(19) World Intellectual Property Organization International Bureau



1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1 (1911) 1

(43) International Publication Date 12 September 2002 (12.09.2002)

PCT

(10) International Publication Number WO 02/070738 A2

(51) International Patent Classification7:

C12Q 1/68

- (21) International Application Number: PCT/CA02/00291
- (22) International Filing Date: 7 March 2002 (07.03.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:

60/273,941 7 March 2001 (07.03.2001) US

- (71) Applicant and
- (72) Inventor: SCHUH, Andre [CA/CA]; 359 Blythwood Road, Toronto, Ontario M4N 1A7 (CA).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): OUWEHAND, Willem [NL/NL]; 147 Hemingford Road, Islington, London N1 1BZ (GB).
- (74) Agent: BERESKIN & PARR; 40 King Street West, 40th Floor, Toronto, Ontario M5H 3Y2 (CA).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

A2

(54) Title: DIAGNOSIS AND TREATMENT OF BLOOD DISORDERS

(57) Abstract: Based on the discovery of the nucleotide and amino acid differences which distinguish the Gov^a and Gov^b allelic forms of the membrane glycoprotein CD109, and which comprise the biallelic Gov platelet alloantigen system, compositions and methods are provided for determining the Gov genotype and phenotype of individuals. Also provided, on the basis of this discovery, are compositions and methods for treating disorders associated with Gov alloantigen incompatibility, such as the bleeding disorders post-transfusion purpura, post-transfusion platelet refractoriness, and neonatal alloimmune thrombocytopenia. The two allelic forms of CD109 differ by a single amino acid. The Gov^a allelic form has Tyr at amino acid position 703 in the CD109 sequence. The Gov^b allelic form has Ser at the same position. This amino acid difference is due to a single change, from A for the Gov^a allele to C for the Gov^b allele, in the CD109 gene.

DIAGNOSIS AND TREATMENT OF BLOOD DISORDERS

FIELD OF THE INVENTION

The present invention provides novel compositions and methods for use in diagnosing the occurrence of certain serious disorders, especially certain bleeding disorders, and novel compositions and methods for use in treating such a disorder, in a person in which the disorder has occurred, and novel compositions and methods for use in avoiding such a disorder, in an individual who is susceptible thereto.

BACKGROUND OF THE INVENTION

Among the disorders, which the invention concerns, are those involving abnormal and excessive bleeding due to destruction of blood platelets ("platelets").

These disorders include, but are not restricted to, post-transfusion purpura ("PTP") and post-transfusion platelet refractoriness ("PTPR"), which are suffered by some persons who receive blood, platelets, leukocyte concentrates, or plasma from other persons by transfusion or the like.

The disorders also include one that is suffered by fetuses and newborns and is known as "neonatal alloimmune thrombocytopenia" ("NATP"). This disorder can cause death of fetuses and serious birth defects or death of newborns. NATP is estimated to affect about 1 in 1000 newborns. In NATP, fetal platelets, which enter the mother's blood stream, induce production in the mother of antibodies directed against fetal platelets. These maternal antibodies then pass with the mother's blood into the fetus and mediate destruction of platelets in the fetus.

A mother, whose fetus or newborn suffers from NATP, is at increased risk of suffering PTP or PTPR.

When platelets from a first human (a "donor") are introduced into the blood system of a second human (a "recipient") by transfusion, through the placenta (in the case of fetal blood entering the mother), or the like, the recipient may mount an immune response against the platelets from the donor. Such an immune response is referred to as an "alloimmune" response, because it involves antibodies reacting against antigens of a different individual of the same species. The alloimmune response to platelets is due to an immune response of the recipient against "alloantigens" (antigens of the same species as that mounting the immune response) on platelets from the donor. These alloantigens are found on membrane glycoproteins that occur in the cell membranes, which define the outer surfaces of platelets ("platelet membranes"). In this invention, the glycoprotein is anchored to the membrane in an atypical manner through an anchor consisting of glycosylphosphatidylinositol (GPI), which anchors an extracellular domain or segment of the glycoprotein exposed to the outside of the platelet. It is thought that

alloantibodies, which are generated in an alloimmune response against platelet alloantig ns, interact with the extracellular domains of the alloantigens.

The platelet alloantigens that a person has are determined by the person's genetics. A donor, because of his or her genetics, may have a platelet alloantigen, which a recipient, who receives blood, platelets, leukocytes or plasma from the donor, does not have, because of the recipient's genetics. In such a situation, the immune system of the recipient may recognize the donor's alloantigen as "non-self," and raise an immune response against, the platelet alloantigen, which the donor has but the recipient does not.

Membrane glycoprotein alloantigens have been characterised for both human red blood cells and human platelets. It is noteworthy, however, that they also occur on other cell types, such as leukocytes and endothelial cells, where they may also occasion various disorders through alloimmune responses.

Recognised classes of red blood cell and platelet alloantigens have been described, over the past 30 years, based on observations of antibody reactions occurring when blood recipients have been exposed to blood from donors.

A recent review of human platelet alloantigen systems is provided by Ouwehand, W., and Navarrete, C., in *Molecular Haematology*, Provan, D. and Gribben, J. eds. Blackwell (1999).

Several biallelic platelet alloantigen "systems" have been characterised. In each of these systems, there are two alloantigens, each of which is provided by one of two alleles of the gene comprising the system. Because each gene occurs twice in the normal human genome, a person can be homozygous for one or the other of the two alloantigens, or heterozygous for the two alloantigens, comprising a biallelic system. The alloantigens described to date occur on glycoprotein molecules which may exist in various forms (transmembrane, GPI-linked and soluble, for example). In such a case, the alloantigens are found on each of the variant forms of the glycoprotein. For all of the biallelic platelet alloantigen systems that have been characterised at the level of protein and gene sequences, it has been found in all cases, except for one, that the difference between the two alleles is based on a single nucleotide polymorphism in the relevant gene.

One biallelic system of human platelet alloantigens is the Gov^a/Gov^b biallelic system associated with CD109, a membrane glycoprotein which occurs on platelets and various other cell types, including leukocytes and endothelial cells. Each Gov allele corresponds to one CD109 glycoprotein (Sutherland, D.R. et al., 1991; Smith et al., 1995; Berry, J. et al., 2000), consistent with the known tissue distribution of CD109. The frequencies for the Gov alleles are 0.4 for Gov^a and 0.6 for Gov^b in the Caucasian population. Thus, in this population, 40.7% are heterozygous for the Gov alleles, and will not mount an alloimmune response due to Gov incompatibility (not possessing the Gov alloantigen found on platelets received from another). In contrast, 19.8% of

Caucasians are homozygous for the Gov^a allele and thus may mount an immune response due to Gov alloantigen incompatibility against platelets received from anyone in the 80.5% of the Caucasian population that is not homozygous for the Gov^a allele, while 39.8% are homozygous for the Gov^b allele and thus may mount an immune response due to Gov alloantigen incompatibility against platelets received from anyone in the 60.2% of the Caucasian population that is not homozygous for the Gov^b allele.

As indicated above, alloimmunization based on Gov incompatibility (the introduction into the blood stream of donor platelets bearing a Gov alloantigen not carried by the recipient) can result in bleeding disorders due to platelet destruction, including NATP, PTPR, and PTP. The location of the Gov antigens within the CD109 molecule, and the nature of the CD109 polymorphism which underlies the Gov^a/Gov^b alloantigen (both at the protein and at the gene level), have not heretofore been known.

Furthermore, it has not heretofore been possible to generate non-human antibody (polyclonal or monoclonal), as from a rat, mouse, goat, chicken, or the like, with specificity for the Gov^a alloantigen but not the Gov^b alloantigen (or vice-versa) sufficient for use in an immunoassay, for typing for Gov phenotype using platelets or CD109 molecules.

Previously developed technology, involving gene-specific amplification of platelet RNA-derived cDNA, followed by the determination of the nucleotide sequence of the amplified DNA, has been applied successfully to the elucidation of the molecular basis of other biallelic platelet alloantigen systems (Newman et al., J. Clin. Invest. 82,739-744 (1988); Newman et al., J. Clin. Invest. 83, 1778-1781 (1989)(P1A or HPA-1 system); Lyman et al., Blood 75, 2343-2348 (1990)(Bak or HPA-3system); Kuijpers et al., J. Clin. Invest. 89, 381-384 (1992)(HPA-2 or Ko system); Wang et al., J. Clin. Invest. 90, 2038-2043 (1992)(Pen system). With one exception, it has been found in each case that a single amino acid difference at a single position differentiates the amino acid sequences of the two alleles, and that this difference arises from a single allele-specific nucleotide substitution in the coding region of the mRNA and gene. There remains a need to elucidate the molecular basis of the biallelic Gov platelet alloantigen system.

SUMMARY OF THE INVENTION

The Gova/Govb CD109 single nucleotide polymorphism

We have now discovered that a single amino acid difference in the CD109 glycoprotein distinguishes the Gov^a and Gov^b allelic forms. The two alleles differ at amino acid position 703 of the full-length 1445 amino acid CD109 molecule, with the Gov^a allele [SEQ ID NO:2] containing a Tyr at this position, while the Gov^b allele [SEQ ID NO:4] contains Ser.

Further, we have discovered that this difference in amino acid sequence between the allelic forms of CD109 is due to a single nucleotide polymorphism at position 2108 of the coding portion of full-length mRNA encoding CD109, or of the corresponding coding strand of the cDNA corresponding to this mRNA. Specifically, the Gov^a allele [SEQ ID NO:1] contains adenine at position 2108, the second nucleotide of the codon encoding the amino acid at position 703 of the full-length CD109 protein, while the Gov^b allele contains cytosine at position 2108, as shown in SEQ ID NO:3

The Gov^a/Gov^b single nucleotide polymorphism of CD109, lies at position 2108 in SEQ ID NO:1. SEQ ID NO:1 is the cDNA sequence encoding the full-length 1445 amino acid CD109 precursor encoding the Gov^a alleleln the Gov^b allele form [SEQ ID NO:3], C occurs at position 2108, rather than A. The ATG at the 5'-end of the sequence in SEQ ID NO:1 corresponds to the translation start of the full-length precursor form (including leader peptide) of CD109. The triplet corresponding to the N-terminal amino acid of the mature CD109 protein is at positions 64-66 in SEQ ID NO:1.

The Gov^a/Gov^b single nucleotide polymorphism of CD109, lies at position 954 in SEQ ID NO:5. SEQ ID NO:5 is the genomic DNA sequence of human CD109 exon 19 and the contiguous introns, introns 18 and 19. The Gov^a/Gov^b single nucleotide polymorphism of CD109 is found within CD109 exon 19, and specifically is located at position 3 of CD109 exon 19. The sequence presented in SEQ ID NO:5 contains A at position 954, and thus corresponds to the Gov^a allele. The corresponding Gov^b sequence contains C at position 954 of SEQ ID NO:5 (nucleotide position 3 of exon 19).

In view of this discovery, it will be readily apparent to the skilled what the present invention provides:

Gov allele-specific oligonucleotides and polynucleotides: Based on the discovery, the present invention provides oligonucleotides and polynucleotides (seems repetitive), including (but not limited to) probes which can be used to determine whether a person is homozygous for one or the other of the Gov alleles, or heterozygous for these alleles, thereby to determine that person's Gov genotype, and by extension, their Gov phenotype (i.e., the Gov alloantigen(s) which their cells express). Further, the invention provides methods of using such oligonucleotides, and test kits to facilitate their use, in such Gov genotype and phenotype determinations. These oligonucleotides of the invention can be used to determine whether, in the CD109 gene, or in the mRNA encoding CD109, the internal nucleotide (nucleotide 2108) of the codon (in CD109 gene or in the mRNA encoding CD109) which corresponds to the amino acid at position 703 in the sequence of full-length CD109 is adenine or cytosine. Such probes will typically be cDNA but may be genomic DNA, mRNA or RNA, and may be labelled for detection. The oligonucleotides of the

invention can be used as probes to detect nucleic acid molecules according to techniques known in the art (for example, see US patent nos. 5,792,851 and 5,851,788).

For example, an oligonucleotide of the invention may be converted to a probe by being end-labelled using digoxigenin-11-deoxyuridine triphosphate. Such probes may be detected immunologically using alkaline-phosphate-conjugated polyclonal sheep antidigoxigenin F(ab) fragments and nitro blue tetrazolium with 5-bromo-4-chloro-3-indoyl phosphate as chromogenic substrate.

Gov allele-specific antibodies: Still further, based on the discovery, which underlies the invention, of the molecular basis for the Gov^a/Gov^b alloantigen system, the invention provides non-human polyclonal and monoclonal antibodies, which can be used to distinguish one Gov allelic form of CD109 from the other, whether the CD109 is part of a complex embedded in or isolated from a membrane or is isolated. These antibodies of the invention, which are preferably provided in an aqueous buffer solution, and the immunoassays of the invention which employ such antibodies, are useful for determining whether a person has one or both of the Gov alloantigens and for Gov phenotyping. Methods of using the antibodies of the invention in the immunoassays of the invention, and in such determinations, are also encompassed by the invention. The invention also provides test kits to facilitate carrying out such immunoassays and determinations.

Gov allele-specific peptides and polypeptides: Again, based on the discovery that underlies the invention, of the molecular basis for the Gov^a/Gov^b alloantigen system, the invention provides peptides or polypeptides, which are useful for various purposes. These peptides or polypeptides are typically between 4 and 100, and more typically between 7 and 50, amino acids in length, and have amino acid sequences identical or having sequence identity to those of segments of the CD109 sequences, that include the amino acid at position 703 of full-length mature CD109. This amino acid (position 703) corresponds the triplet at positions 2107-2109 in the CD109 cDNA sequence presented in SEQ ID NO:1, or in the corresponding sequence for the CD109 cDNA that encodes the Gov^b allelic form [SEQ ID NO:3]. These peptides or polypeptides may be synthetic, may be purified from native CD109 or may be prepared by recombinant means. For quidance, one may consult the following US patent nos. 5,840,537, 5,850,025, 5,858,719, 5,710,018, 5,792,851, 5,851,788, 5,759,788, 5,840,530, 5,789,202, 5,871,983, 5,821,096, 5,876,991, 5,422,108, 5,612,191, 5,804,693, 5,847,258, 5,880,328, 5,767,369, 5,756,684, 5,750,652, 5,824,864, 5,763,211, 5,767,375, 5,750,848, 5,859,337, 5,563,246, 5,346,815, and WO9713843. Many of these patents also provide guidance with respect to experimental assays, probes and antibodies, methods, transformation of host cells, which are described below. These patents, like all other patents, publications (such as articles and database publications) in this application, are incorporated by reference in their ntirety.

Gov allele-sp cific peptides and polypeptides as antig ns and immunogens, and Gov allel -sp cific p lycl nal and monoclonal antib dies: These peptides or polypeptides are useful as antigens (usually coupled to a larger, immunogenic carrier [proteinaceous or otherwise], as known in the art) for making the polyclonal or monoclonal antibodies of the invention. The peptides or polypeptides are also useful in screening monoclonal antibody-producing cultures (hybridoma cultures/E. coli cultures or so-called V gene phage antibodies) to identify those that produce monoclonal antibodies of the invention.

The invention also encompasses immunogenic compositions which comprise a peptide, polypeptide or fusion compound of the invention and which are immunogenic in a bird, including, without limitation, a chicken, or a mammal, such as, a mouse, rat, goat, rabbit, guinea pig, sheep or human. The compositions may include an immunogenicity-imparting "carrier" which may be but is not necessarily a protein as known in the art, that is immunogenic in a bird or mammal, coupled to at least one peptide or polypeptide of the invention, which has an amino acid sequence that is the same as that of a segment of the sequence for CD109, that includes the amino acid at position 703 of the full length CD109 molecule.

The present invention also provides methods of using the peptides, polypeptides and immunogenic compositions of the invention for making antibodies of the invention, and methods of using the peptides and polypeptides of the invention in screening monoclonal antibody-producing hybridoma cultures or bacterial clones for those that produce monoclonal antibodies or fragments thereof of the invention.

Therapeutic and diagnostic application of Gov allele-specific peptides, polypeptides, and antibodies: These peptides or polypeptides, as well as antibodies, which are specific for the Gov^a [SEQ ID NO:2] or Gov^b [SEQ ID NO:4], but not both, allelic forms of CD109 in the platelet membrane, and which can be produced by a mammal (including an human) immunized with the peptides or polypeptides, which themselves happen to be immunogenic, or the immunogenic compositions of the invention, are also useful both therapeutically and diagnostically. The invention also provides the methods of using the peptides and polypeptides of the invention, and antibodies made using the peptides that are immunogenic and the immunogenic compositions of the invention, in therapeutic and diagnostic applications.

The Gov allele-specific peptides or polypeptides can also be used diagnostically to detect the presence of Gov^a or Gov^b specific antibodies in human plasma or serum samples, using methods that are readily apparent to those skilled in the art. Such analyses would be useful in the investigation of cases of acquired alloimmune thrombocytopenia, including PTP, PTPR, and NATP. In the latter case, this approach could also be used to detect the presence of Gov allele-specific antibodies in the mother of the affected fetus or newborn. The presence of Gov allele-

specific antibodies can also be detected using platelets of known Gov phenotype. However, this approach has numerous technical disadvantages that are eliminated by the use of Gov allelespecific peptides or polypeptides for Gov allelespecific antibody detection.

Administration to a person, who is suffering from, or at risk for, for example, PTP or PTPR, or a mother at risk for passing NATP-causing alloantibodies to her fetus, of one of the peptides or polypeptides, that would be bound by the anti-Gov alloantibodies in such a person, would inhibit the binding of the alloantibodies to the person's (or the fetus's platelets and thereby inhibit the platelet destruction and abnormal bleeding associated with the disorders. Alternatively, administration to such a person of antibodies (particularly human antibodies), which are produced using a peptide or polypeptide of the invention, which is immunogenic by itself, or an immunogenic composition of the invention, and which are specific for the Gov allelic form of the CD109 on the person's platelets which is associated with the PTP or PTPR, from which the person is suffering or may suffer, would induce the production of anti-idiotypic antibodies, which, in turn, would inhibit the platelet-destructive effects of the anti-Gov alloantibodies, which are generated by the person's own immune system and which are causing or threatening to cause the PTP, PTPR or NATP. These therapeutic applications of peptides and polypeptides of the invention would be especially useful in treating NATP in a newborn, because the alloantibody giving rise to NATP in the newborn is not continuously produced by the immune system of the newborn, but rather is acquired passively, and therefore in limited, non-replenished quantity, by the newborn from its mother.

Thus, in accordance with one aspect of the present invention, an oligonucleotideprobe is provided that hybridizes to a portion of the CD109 gene, or a portion of CD109-encoding mRNA or cDNA prepared from such mRNA, which portion includes a nucleotide corresponding to the internal nucleotide of the codon for the amino acid at position 703 of the full-length CD109 molecule, and that is capable of distinguishing one Gov allele from the other through the ability to hybridize under stringent conditions to the portion in question only when the nucleotide in question is A (or dA), when the probe is to detect the Gov^a allele, or C (or dC), when the probe is to detect the Gov^b allele. The nucleotide in question is at position 2108 of the coding region of the CD109 cDNA sequence and lies at position 2108 in SEQ ID NO:1. The cDNA sequence has A at this position, and so is the sequence corresponding to the Gov^a allele. The nucleotide in question lies at position 954 of the sequence presented as SEQ ID NO:5 and contains an A in this position, and thus also corresponds to Gov^a allele.

The Gov allele-specific oligonucleotide hybridization probes of the invention may comprise genomic DNA, cDNA, or RNA, although preferably it is DNA. Such oligonucleotide probes can be synthesised by automated synthesis and will preferably contain about 10 - 30 bases, although as understood in the oligonucleotide probe hybridization assay art, as few as 8

and as many as about 50 nucleotides may be useful, depending on the position within the probe where the potential mismatch with the target is located, the extent to which a label on the probe might interfere with hybridization, and the physical conditions (e.g., temperature, pH, ionic strength) under which the hybridization of probe with target is carried out.

In accordance with another aspect of the present invention, a test kit for Gov alloantigen typing is provided comprising:

(a) means for amplifying nucleic acid that comprises at least a portion of a CD109 gene, a CD109-encoding mRNA, or a CD109 cDNA made from such RNA, wherein the portion includes a nucleotide (nucleotide 2108 in SEQ ID NO:1, or nucleotide 954 in SEQ ID NO:5) corresponding to the internal nucleotide of the codon encoding amino acid 703 of the full length CD109 protein. (b) an oligonucleotide probe of the invention, that distinguishes one Gov allele from the other. The "means for amplifying" will, as the skilled will readily understand, depend on the amplification method to be used. Thus, for example, these means might include suitable primers, a suitable DNA polymerase, and the four 2'-deoxyribonucleoside triphosphates (dA, dC, dG, dT), if amplification is to be by the PCR method. To cite another example, if the amplification is to be by a method relying on transcription, such as the 3SR method, the means will include two primers, at least one of which, when made double-stranded, will provide a promoter, an RNA polymerase capable of transcribing from that promoter, a reverse transcriptase to function in primer-initiated, DNA-directed and RNA-directed, DNA polymerization and possibly also in RNAse H degradation of RNA to free DNA strands from RNA/RNA hybrids, the four ribonucleoside triphosphates (A, C, G and U), and the four 2'-deoxyribonucleoside triphosphates. In another example, if the amplification is by the ligase chain reaction, the means will include two oligonucleotides (DNAs) and a suitable DNA ligase that will join the two if a target, to which both can hybridize adjacent to one another in ligatable orientation, is present.

The oligonucleotide probes of the invention will preferably be labelled. The label may be any of the various labels available in the art for such probes, including, but not limited to ³²P; ³⁵S; biotin (to which a signal generating moiety, bound to or complexed with avidin can be complexed); a fluorescent moiety; an enzyme such as alkaline phosphatase (which is capable of catalysing a chromogenic reaction); digoxigenin, as described above; or the like.

As indicated in the examples, RFLP analysis can be employed, using BstNI (or isoschizomers thereof), in analysing cDNA or genomic DNA (with or without amplification) to determine Gov genotype. As indicated further in the examples, electrophoretic SSCP analysis may be used to determine Gov genotype. And as indicated in the examples, the hybridization studies outlined above may use fluorescent probes, and may be directly coupled to the DNA amplification step, as in "Real-Time PCR" or related methods.

There has also been provided, in accordance with another aspect of the present invention, a method of typing for Gov allele-specific target sequence in a CD 109 nucleic acid derived from a subject, comprising the steps of,

- (a) obtaining, by a target nucleic acid amplification process applied to mRNA from human platelets, endothelial cells, or T cells, an assayable quantity of amplified nucleic acid with a sequence that is that of a subsequence (or the complement of a subsequence) of the mRNA that encodes a CD109 said subsequence including the nucleotide at the position in the mRNA corresponding to position 2108 in SEQ ID NO:1 or to nucleotide 954 in SEQ ID NO:5; and
- analyzing (e.g., in a nucleic acid probe hybridization assay employing an oligonucleotide (b) probe or probes according to the invention) the amplified nucleic acid obtained in step (a) to determine the base or bases at the position in the amplified nucleic acid that corresponds to position 2108 in SEQ ID NO:1 or to nucleotide 954 in SEQ ID NO:5. It is noteworthy that, if the product of the amplification is double-stranded DNA, analysis for Gov genotype can be carried out by a RFLP (restriction fragment length polymorphism) analysis comprising exposing the amplified DNA to the restriction endonuclease BstNI (or isoschizomer thereof) under conditions whereby the DNA will be cleaved if it includes a site for cleavage by that enzyme. Such DNA, prepared from mRNA encoding the Gov^b alloantigen, containing a C rather than an A at the position corresponding to nucleotide 2108 in SEQ ID NO:1 (or to nucleotide 954 in SEQ ID NO:5), includes a recognition site for that endonuclease, while such DNA prepared from mRNA encoding the Goy^a alloantigen, does not. If the analysis, by whatever method, of the amplified nucleic acid reveals that there is only an A (or dA) at the position corresponding to position 12108, the platelets (and blood from which they came) have only the Gov^a alloantigen, and the individual from whom the platelets came, is homozygous for Gov^a. Alternatively, if the analysis of the amplified nucleic acid reveals that there is only a C (or dC) at the position corresponding to position 2108, the platelets (and blood from which they came) have only the Gov^b alloantiden and the individual, from whom the platelets came, is homozygous for the Govb allele. Finally, if the analysis indicates that there is either an A (or dA) or a C (or dC) at that position, the platelets (and blood from which they came) have both Gov alloantigens, and the individual from whom the platelets came, is heterozygous for Gov alloantigen.

In one application of the typing methods of the invention, the methods are applied to two individuals to determine whether blood or platelets from one would provoke an alloimmune response, and possibly PTP or PTPR, in the other. The typing method can be applied with a man and a woman, who are contemplating conceiving or have conceived a child together, to determine the risk that the child would be at risk for NATP and the risk that the woman would be at increased risk for PTP or PTPR. If the woman were heterozygous for the Gov alloantigens there would be, due to Gov alloantigen incompatibility, no risk of NATP and no increased risk for

the woman of PTP or PTPR. If, however, the woman were homozygous for one of the Gov alloantigens, there would be, due to Gov alloantigen incompatibility, risk of NATP in a child and increased risk of PTP or PTPR for the woman, unless the man is homozygous for the same Gov alloantigen as is the woman.

In accordance with yet another aspect of the present invention, a method of typing an individual for Gov alloantigen is provided that comprises analyzing the genomic DNA of the individual to determine the Gov alloantigen(s) of the individual. Applications of this method are substantially the same as those of the method of the invention for typing for Gov alloantigen that begins with platelet, endothelial cell, or T cell mRNA.

This method of the invention, entailing analysis of genomic DNA, can be carried out in substantially the same way as outlined above for analysis of mRNA, namely first amplifying the genomic DNA and then analyzing to product of the amplification to ascertain whether there is only dA, only dC, or both dA and dC, at the position in the coding region of the genomic DNA corresponding to position 2108 in SEQ ID NO:1, or to nucleotide 954 in SEQ ID NO:5.

In accordance with a further aspect of the present invention, a test kit for Gov alloantigen typing is provided comprising a non-human antibody (or antibodies) that distinguishes the two allelic forms of CD109. The antibody (or antibodies) of the kit may be polyclonal, or preferably monoclonal, and in addition to its (their) specificity for either but not both Gov alloantigens (on the surface of platelets or separated therefrom) or the CD109 subunit of one but not both of such alloantigens, typically will recognise a polypeptide molecule encoded by a nucleotide sequence encoding at least amino acid 703 of a CD109 polypeptide (the amino acid at the position corresponding to nucleotides 2107 - 2109 in SEQ ID NO:1, or to nucleotides 953 – 955 in SEQ ID NO:5).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The following definitions are provided in order to aid those skilled in the art in understanding the detailed description of the present invention.

The term "alloantigens" refers to antigens of an individual that are responsible for eliciting an alloimmune response.

The phrase "alloimmune response" refers to an immune response, which occurs when antibodies from one individual react against antigens of a different individual of the same species.

The phrase "anti-idiotypic antibodies" refers to antibodies which can bind endogenous or foreign idiotypic antibodies and which can be used to treat or prevent pathological conditions associated with an immune response to a foreign alloantigen.

The phrase "Gov^a/Gov^b biallelic system" refers to a system of human platelet alloantigens in which an individual can be homozygous for either Gov^a or Gov^b allelic forms of CD 109, or an

individual can be Gov^a/Gov^b heterozygous for CD109.

"GPI" refers to glycosylphosphatidylinositol.

The term "NATP" refers to neonatal alloimmune thrombocytopenia.

"Nucleic acid" includes DNA and RNA, whether single or double stranded. The term is also intended to include a strand that is a mixture of nucleic acids and nucleic acid analogs and/or nucleotide analogs, or that is made entirely of nucleic acid analogs and/or nucleotide analogs.

"Nucleic acid analogue" refers to modified nucleic acids or species unrelated to nucleic acids that are capable of providing selective binding to nucleic acids or other nucleic acid analogues. As used herein, the term "nucleotide analogues" includes nucleic acids where the internucleotide phosphodiester bond of DNA or RNA is modified to enhance biostability of the oligomer and "tune" the selectivity/specificity for target molecules (Ulhmann, et al., 1990, Angew. Chem. Int. Ed. Eng., 90: 543; Goodchild, 1990, J. Bioconjugate Chem., I: 165; Englisch et al., 1991, Angew, Chem. Int. Ed. Eng., 30: 613). Such modifications may include and are not limited to phosphorothloates, phosphorodithioates, phosphotriesters, phosphoramidates or methylphosphonates. The 2'-O-methyl, allyl and 2'-deoxy-2'-fluoro RNA analogs, when incorporated into an oligomer show increased biostability and stabilization of the RNA/DNA duplex (Lesnik et al., 1993, Biochemistry, 32: 7832). As used herein, the term "nucleic acid analogues" also include alpha anomers (α-DNA), L-DNA (mirror image DNA), 2'-5' linked RNA, branched DNA/RNA or chimeras of natural DNA or RNA and the abovemodified nucleic acids. For the puposes of the present invention, any nucleic acid containing a "nucleotide analogue" shall be considered as a nucleic acid analogue. Backbone replaced nucleic acid analogues can also be adapted to for use as immobilised selective moieties of the present invention. For purposes of the present invention, the peptide nucleic acids (PNAs) (Nielsen et al., 1993, Anti-Cancer Drug Design, 8: 53; Engels et al., 1992, Angew, Chem. Int. Ed. Eng., 31: 1008) and carbamate-bridged morpholino-type oligonucleotide analogs (Burger, D.R., 1993, J. Clinical Immunoassay, 16: 224; Uhlmann, et al., 1993, Methods in Molecular Biology, 20,. "Protocols for Oligonucleotides and Analogs," ed. Sudhir Agarwal, Humana Press, NJ, U.S.A., pp. 335-389) are also embraced by the term "nucleic acid analogues". Both exhibit sequence-specific binding to DNA with the resulting duplexes being more thermally stable than the natural DNA/DNA duplex. Other backbone-replaced nucleic acids are well known to those skilled in the art and may also be used in the present invention (See e.g., Uhlmann et al 1993, Methods in Molecular Biology, 20, "Protocols for Oligonucleotides and Analogs," ed. Sudhir Agrawal, Humana Press, NJ, U.S.A., pp. 335).

The term "PTP" refers to post-transfusion purpura.

The term "PTPR" refers to post-transfusion platelet refractorines.

"SNP" refers to single nucleotide polymorphism.

Th standard, one-letter codes "A," "C," "G," and "T" are used herein for the nucleotides adenylate, cytidylate, guanylate, and thymidylate, respectively. The skilled will understand that, in DNAs, the nucleotides are 2'-deoxyribonucleotide-5'-phosphates (or, at the 5'-end, possibly triphosphates) while, in RNAs, the nucleotides are ribonucleotide-5'-phosphates (or, at the 5'-end, possibly triphosphates) and uridylate (U) occurs in place of T. "N" means any one of the four nucleotides. On occasion herein, dA, dC, dG and dT might be used for the respective 2'-deoxyribonucleotides.

Unless otherwise specified or required by the context, "nucleic acid" means DNA or RNA and "nucleotide" means ribonucleotide or 2'-deoxyribonucleotide.

Reference herein to a "full-length" CD109 molecule or protein means the 1445-amino acid-long polypeptide, for which the amino acid sequence, deduced from a cDNA sequence, is provided in SEQ ID NO:1 and in SEQ ID NO:3 and which is denoted as the full-length translated product (i.e., including the amino-terminal leader peptide, and excluding carboxyl-terminal processing associated with GPI anchor addition). The Gov^a alloantigen bearing form of CD109 may be referred to herein as ⁷⁰³Tyr CD109. The Gov^b alloantigen bearing form of CD109 may be referred to herein as ⁷⁰³Ser CD109.

It has been determined that a single nucleotide of the CD109 gene is responsible for the Gov polymorphism in CD109. Extensive serological studies initially demonstrated that the polymorphism underlying the Gov system resides solely on the CD109 molecule [Sutherland, D.R. (1991); Smith et al. (1995)]. Further, extensive deglycosylation of CD109 does not affect the binding the anti-Gov^a and anti-Gov^b antibodies to molecules of the appropriate phenotype, or to cells bearing the appropriate CD109 variant, indicating that carbohydrate residues are not involved in the formation of Gov antigenic epitopes. Further work has indicated that the Gov allele-specific antibody binding can however, be abrogated by denaturation of CD109 with the detergent SDS [Smith et al. (1995)]. Taken together, these observations indicate that the Gov alleles of CD109 are protein epitopes that are likely defined by the primary amino acid sequence of CD109.

Following the isolation of a CD109 cDNA the nature of the two Gov alleles was characterised further using platelet RNA-derived cDNA in the polymerase chain reaction ("PCR"). Platelet mRNA transcripts were obtained from serologically defined Gov^{a/a}, Gov^{a/b} and Gov^{b/b} individuals. The RNA was then converted to cDNA, and the entire CD109 cDNA coding region was then amplified as a series of overlapping PCR products. The Gov^a [SEQ ID NO:1] and Gov^b [SEQ ID NO:3] alleles differ by an A to C substitution at position 2108 of the coding region of the CD109 cDNA. This single nucleotide polymorphism also results in a BstNI restriction site in the Gov^b allele that is not present in its Gov^a counterpart. On the basis of this BstNI site, Gov^a can

by distinguished from Gov^b by restriction fragment length polymorphism (RFLP) analysis. This single nucleotide polymorphism can also be detected by SSCP analysis, and by allele-specific hybridization studies, including "Real-Time" PCR analyses.

As a result of this A²¹⁰⁸C single nucleotide polymorphism, the Gov^a all le [SEQ ID NO:2] of CD109 contains a Tyr at position 703 of the full-length protein, while the Gov^b allele [SEQ ID NO:4] contains a Ser in this position. The polymorphism does not alter the ability of Gov^a and Gov^b homozygous platelets to adhere to collagen types I, III and V. Additionally, the binding of anti-Gov^a and anti-Gov^b antibodies to platelets of the appropriate phenotype did not interfere with platelet adhesion to any of the above collagen types. Thus, while the Tyr⁷⁰³Ser results in the formation of the Gov alloantigen epitopes, it does not appear to impair platelet function.

Identification and characterisation of the Gov alloantigen system permits pre- and post-natal diagnosis of the Gov phenotype of an individual, providing a warning for the possibility of NATP, PTP and PTPR. Allelic Gov typing of CD109 equates with the Gov status of the CD109 protein of an individual. The Gov system led to diagnostic and therapeutic strategies to avoid or control diseases that result from Gov incompatibility. The present invention can be applied to these tasks and goals in a variety of ways, illustrative examples of which are discussed below.

For example, an oligonucleotide probe can be synthesized, in accordance with the present invention, that will hybridize to a cDNA segment, derived from CD109 mRNA, that contains the nucleotide G at polymorphic nucleotide 2108 (nucleotide=guanylate). Alternatively, an oligonucleotide probe can be synthesized that will hybridize with a CD109 cDNA segment containing the base adenine at nucleotide 2108. (nucleotide=adenylate). These allele-specific probes can be appropriately labelled and added to the generated cDNA segments under annealing conditions, such that only one of the allele-specific probes hybridizes and can be detected, thereby identifying the specific Gov^a or Gov^b allele. In accordance with conventional procedures, the design of an oligonucleotide probe according to the present invention preferably involves adjusting probe length to accommodate hybridization conditions (temperature, ionic strength, exposure time) while assuring allele-specificity. A length of ten to thirty nucleotides is typical.

Diagnostic kits can also be used, in accordance with the present invention, for the determination and diagnosis of alloantigen phenotypes via the procedures described herein. Such a kit can include, among others, antibodies or antibody fragments to an antigenic determinant expressed by either of the above-described Gov^a- and Gov^b- encoding sequences. These antibodies would react with the blood sample of an individual so as to indicate whether that individual has a Gov^a or Gov^b phenotype. Alternatively, all the reagents required for the detection of nucleotide(s) that distinguish the Gov alloantigens, by means described herein, can be provided in a single kit that us a isolated genomic DNA, platelet (or other cellular) mRNA or

total RNA, or corresponding cDNA from an individual. A kit containing a labelled probe that distinguishes, for example, nucleotide 2108 of CD109 can be utilised for Gov alloantigen genotyping and phenotyping.

A further beneficial use of the nucleotide sequences that distinguish the Gov^a allele from the Gov^b allele is to obtain or synthesize the respective expression product, in the form of a peptide or polypeptide, encoded by these nucleotide sequences. These polypeptides can be used to generate antibodies for diagnostic and therapeutic uses, for example, with regard to pathological conditions such as PTP, PTPR or NATP. These polypeptides can also be used diagnostically to detect the presence of Gov^a or Gov^b specific antibodies in patient plasma or serum, or used therapeutically (see below; assays may be adopted, for example, from US Patent No. 5,851,788).

A polypeptide within the present invention which can be used for the purpose of generating such antibodies preferably comprises an amino-acid sequence that corresponds to (i.e., is coincident with or functionally equivalent to) a fragment of the CD109 molecule that includes amino acid 703. When amino acid 703 is Tyrosine, the polypeptide can be used, as described above, to produce antibodies that specifically bind the Gov^a form of CD109; in contrast, when it is Serine, antibodies can be obtained that specifically recognise the Gov^b form. The class of polypeptides thus defined, in accordance with the present invention, is not intended to include the native CD109 molecule, but does encompass fragments of the molecule, as well as synthetic polypeptides meeting the aforementioned definition.

Although the length of a polypeptide within this class is not critical, the requirement for immunogenicity may require that the polypeptide be attached to an immunogenicity-imparting carrier. Such carriers include a particulate carrier such as a liposome or a soluble macromolecule (protein or polysaccharide) with a molecular weight in the range of about 10,000 to 1,000,000 Daltons Additionally, it may be desirable to administer the polypeptide with an adjuvant, such as complete Freund's adjuvant. For artificial polypeptides, as distinguished from CD109 fragments, maximum length is determined largely by the limits of techniques available for peptide synthesis, which are currently about fifty amino acids. Thus, a synthetic polypeptide of the present invention is preferably between four to about fifty amino acids in length.

In the context of the present invention, the term "antibody" encompasses monoclonal and polyclonal antibodies produced by any available means. Such antibodies can belong to any antibody class (IgG, IgM, IgA, etc.) and may be chimeric. Examples of the preparation and uses of polyclonal antibodies are disclosed in U.S. Patent Nos. 5,512,282, 4,828,985, 5,225,331 and 5,124,147 which are incorporated by reference in their entirety The term "antibody" also encompasses antibody fragments, such as Fab and F(ab')₂ fragments, of anti-Gov^a or anti-Gov^b antibodies, conjugates of such fragments, and so-called "antigen binding proteins" (single-chain

antibodies) which are based on anti-Gov^a or anti-Gov^b antibodies, in accordance, for example, with U.S. Pat. No. 4,704,692, the contents of which are hereby incorporated by reference. Alternatively, monoclonal antibodies or fragments thereof within the present invention can be produced using conventional procedures via the expression of isolated DNA that encodes variable regions of such a monoclonal antibody in host cells such as E. coli (see, e.g., Ward et al., Nature, 341:544-546 (1989)) or transfected murine myeloma cells (see Gillies et al., Biotechnol. 7:799-804 (1989); Nakatani et al., Biotechnol. 7:805-810 (1989)). For additional examples of methods of the preparation and uses of monoclonal antibodies, see U.S. Patent Nos. 5,688,681, 5,688,657, 5,683,693, 5,667,781, 5,665,356, 5,591,628, 5,510,241, 5,503,987, 5,501,988, 5,500,345 and 5,496,705 that are incorporated by reference in their entirety.

While human alloantisera currently used for serological typing are specifically excluded from this definition, the use of CD109 or Gov allele-specific peptides to detect anti-Gov antibodies in human plasma or serum, or to determine the specificity of such alloantibodies, are specifically included. Similarly, the use of such CD109 peptides or Gov allele-specific peptides to purify CD109 antibodies, or allele-specific CD109 antibodies from human serum is specifically included. Similarly, the use in vitro of such CD109 peptides or Gov allele-specific peptides to deplete allele-specific antibody activity from human serum samples, or to block CD109 antibody binding, or allele-specific antibody binding, is specifically included.

Diagnostic applications of these antibodies are exemplified, according to the present invention, by the use of a kit containing an anti-Gov^a or an anti-Gov^b antibody, which undergoes a reaction with a sample of an individual's blood to determine a Gov^a or Gov^b platelet phenotype. Such a reaction involves the binding of anti-Gov^a antibody to Gov^a antigen or the binding of anti-Gov^b antibody to Gov^b antigen. The observation of antibody-antigen complex in a blood sample would indicate a positive result. A kit of this type could be used to diagnose, or to help prevent the occurrence of pathological conditions like PTP, PTPR, or NATP.

A polypeptide of the present invention that is recognised specifically by anti-Gov^a or anti-Gov^b antibodies can also be used therapeutically. Thus, antibodies raised against such a polypeptide can be employed in the generation, via conventional methods, of anti-idiotypic antibodies, that is, antibodies that bind an anti-Gov^a or anti-Gov^b antibody. See, e.g., U.S. Pat. No. 4,699,880, the contents of which are hereby incorporated by reference. Such anti-idiotypic antibodies would bind endogenous or foreign anti-Gov antibodies in the blood of an individual, which would treat or prevent pathological conditions associated with an immune response to a "foreign" Gov alloantigen. Alternatively, a polypeptide within the present invention can be administered to an individual, with a physiologically-compatible carrier, to achieve the same qualitative effect, namely, the selective reduction or elimination of circulating anti-Gov antibodies from a patient suffering or at risk from an immune response, or the abrogation by competitive

binding to administered peptide, of the binding of Gov-specific antibodies to the platelets of such an individual

The present invention is further described below by reference to the following, illustrative examples.

EXAMPLE 1: PCR Amplification and Analysis of PCR Products

Platelet total RNA was isolated from EDTA anticoagulated blood of Gov^{aa} and Gov^{bb} individuals in the manner described in Lymann et al., Blood 75:2343-48 (1990). First, platelet mRNA in 10 μl aliquots was heated to 70°C for 10 minutes and quickly cooled on ice before reverse transcription. The first strand cDNA was then synthesized using 10 μM oligo dT, 40 units RNAsin (Promega), 2 mM of each dNTP (dN triphosphate) (Pharmacia), 500 units of cloned MMLV reverse transcriptase and 5x enzyme buffer (Gibco) in a total volume of 50 μl. The cDNA synthesis was carried out at 42°C for 45 minutes and was stopped by chilling to 0°C.

Overlapping sets of oligonucleotide primers (Table 2) based on the sequence of CD109 were then used to amplify by PCR the entire coding region of platelet CD109 in 8 overlapping segments that spanned the entire CD109 open reading frame.

TABLE 2:

Fragm ent	Sense Primer		Antisense Primer	-	Size (bp)	Annes ng Tempi ature (°C)
1	K1-80 5' GTAGCCCAGGCAGACGCC 3'	(-24)	K1-650 5' GTGACAACCACTGTTGGATCAA 3'	544	568	59
2	K1-1 5' CGCATTGTTACACTCTTCTC 3'	445	K1-1120 5' TACATTTCTTGAAATACCTG 3'	1014	570	50
3	K1-1022 5' GATTCTTCAAATGGACTTT 3'	910	K1-REV-1 5' GGCTGTGTCACAGAGATC 3'	1747	838	50
4	K1-1400 5' TGAATTCCCAATCCTGGAGGA 3'	1291	GSP3 5' GCCACCCAAGAAGTGATAGA 3'	2165	875	55
5	K1-M43 5' TTCAGGAATGTGGACTCTGG 3'	1898	6R4N 5' CGGCTTCAAGGAAACATCT 3'	2998	1101	56

. 6	K1-3080 5' CTGGGAGCACTTGGTTGTCA 3'	2948	1-5N 5' CAGCAACATCTAAATCAAAGGC 3'	3859	912	56
7	K1-3570 5' ACAATTTCAGACTTCTGAGG 3'	3462	7U3N 5' CACAGCCAAAGTTCCATA 3'	4337	876	50
8	K1-3920 5' GACGAAGATCTATCCAAAATC 3'	3812	K1-4600 5' GCTAGGACCTGTTGTACACC 3'	4489	678	55

Table 2 lists the position of the 5' end of each oligonucleotide with respect to the CD109 cDNA sequence, which includes both 3' and 5' untranslated regions, is noted in parentheses. The CD109 ORF encompasses nucleotides 1-4335 of the published CD109 cDNA, and corresponds exactly to the CD109 cDNA sequence presented in SEQ ID NO:1. The size of each PCR product, and the annealing temperature used for the corresponding primer pair, is listed.

PCR reactions (50:1) containing 1x PCR buffer (Gibco Life Technologies), 1.5 mM MgCl₂, 200 :M of each dNTP, 1 :M of each primer, 1.25 units Taq polymerase (Gibco Life Technologies), and 3: I cDNA underwent 40 cycles of 94°C (45 seconds), primer-specific annealing temperature (Table 2: 45 seconds), and 72°C (45 - 60 seconds), using a Perkin Elmer 2400 thermocycler. PCR products (30 :I) were subsequently size-separated electrophoretically on a 1.2 % agarose/TAE gel containing 1:g/ml ethidium bromide. Bands were subsequently excised and purified (50:1) using the QIAquick (Qiagen) kit for direct sequencing and subcloning. Sequencing reactions (3-5: purified product per reaction) were carried out using the Thermosequenase Cy5.5 dye terminator sequencing kit (Amersham Pharmacia Biotech) and the same primers that had been used for initial PCR amplification (Table 2), or selected internal CD109-specific primers as appropriate, and were subsequently analysed using the Open Gene automated DNA sequencing system (Visible Genetics). In parallel, PCR products were cloned into Pmel-digested pMAB1, a pBS SK(-) (Stratagene) derivative containing a Pmel restriction site within the polylinker. Resultant plasmid clones were analysed by alkaline lysis/restriction digestion, and as appropriate (and following an additional overnight 13% PEG/ 1.6 M NaCl precipitation), by DNA sequence analysis as above. By combining direct PCR sequencing and the analysis of subcloned fragments, it was ensured that the DNA sequence of each PCR-derived cDNA fragment was obtained independently at least twice, with each fragment being sequenced in both directions in its entirety.

This analysis revealed that the CD109 cDNA sequences of Gov^{aa} and Gov^{bb} individuals differed by a single nucl otide at position 2108 of the sequence shown in SEQ ID NO:1. Gov^{a/a}

individuals have an A at position 2108, whereas Gov bib individuals have a C at the same position. This change results in a Tyr-Ser amino acid polymorphism at residue 703 of the full-length CD109 polypeptide chain. This single nucleotide polymorphism also results in a BstNI restriction site in the Govb allele that is not present in the Gov^a allele. Analysis of the other regions of the CD109 cDNA in their entirety revealed no other nucleotide differences that segregated with Gov phenotype (i.e., that could be used to distinguish the Gov^a allele from the Gov^b allele).

To facilitate subsequent genomic DNA analyses of the *Gov*^{ab} alleles, the intron/exon junctions of the exon bearing the putative Gov-specific nucleotide substitution identified above, as well as the DNA sequence of the flanking introns, were determined. CD109 cDNA-specific oligonucleotides binding in the vicinity of this substitution were used for the direct sequencing of p4L10, a pCYPAC_1-derived PAC clone bearing the human CD109 locus using the Open Gene system (Visible Genetics) as above. The nucleotide sequence of the Gov polymorphism-containing exon, as well as of the flanking introns, is presented in SEQ ID NO:5. The Gov polymorphism lies at nucleotide position 954 in SEQ ID NO:5. Subsequent work has mapped the intron-exon structure of the entire human CD109 locus, and has determined that the Gov single nucleotide polymorphism of CD109 lies in exon 19 of the CD109 gene.

EXAMPLE 2: RFLP Analysis of PCR Amplified Genomic DNA

The A-C Gov CD109 polymorphism corresponds to the internal nucleotide of the first complete codon of exon 19 of the CD109 gene. As this exon comprises only 118 nucleotides, and the Gov polymorphism lies almost at the extreme 5' end of this exon, we determined the nucleotide sequence of both introns flanking this exon to facilitate subsequent genomic DNA analyses of the Gov^{a/b} alleles. The DNA sequence of CD109 exon 19 and its flanking introns (CD109 introns 18 and 19) is presented as SEQ ID NO:5. To confirm that the A to C polymorphism at position 2108 of the CD109 open reading frame (nucleotide 2108, SEQ ID NO: 1; nucleotide 954, SEQ ID NO:5) segregates with the Gov phenotype, RFLP analysis was carried out on PCR amplified genomic CD109 DNA using the BstNl restriction endonuclease, which recognises the DNA sequence 5' CCAGG 3' found in the Gov^b cDNA (nucleotides position 2108 - 2112 in SEQ ID NO:3; the corresponding Gov^a sequence, 5' ACAGG 3', is nucleotides 2108 - 2112 in SEQ ID NO:1). This enzyme does not cleave at 5' ACAGG 3' (found in Gova; nucleotides 2108 - 2112 in SEQ ID NO: 1), A 448 bp genomic fragment was PCR-amplified from . Gov^{aa}, Gov^{ab}, and Gov^{bb} individuals using the pair of oligonucleotides SEQ ID NO:9 and SEQ ID NO:10. These oligonucleotides flank exon 19. The former binds within intron 18 (nucleotides 875). - 892 SEQ ID NO:5), while the latter binds within intron 19 to the sequence complementary to nucleotides 1305 - 1322 of SEQ ID NO:5). The resultant 448 bp PCR product, when digested

with BstNI, yielded the restriction fragments predicted on the basis that the A to C polymorphism at position 2108 (SEQ ID NO: 1) segregates with the Gov phenotype.

EXAMPLE 3: Hybridization Analysis of PCR Amplified Genomic DNA

To further confirm that the A to C polymorphism at position 2108 of the CD109 open reading frame (nucleotide 2108, SEQ ID NO:1; nucleotide 954, SEQ ID NO:5) segregates with the Gov phenotype, we also performed an alternative analysis involving the selective hybridization of Gov allele-specific DNA probes to PCR amplified genomic CD109 DNA. Two primers flanking the polymorphic A-C site at position 2108 (SEQ ID NO:1; position 954, SEQ ID NO:5) were designed to amplify by PCR a 105 bp genomic DNA fragment containing the polymorphic site from genomic DNA isolated from Gov^{aa}, Gov^{ab}, and Gov^{bb} individuals. The first primer (SEQ ID NO:11) binds within intron 18 to nucleotides 902 – 928 of SEQ ID NO:5. The second primer (SEQ ID NO:12) binds within exon 19 to the sequence complementary to nucleotides 977 - 1106 of SEQ ID NO:5. Two additional nucleotide probes were designed - one specific for the target sequence of the Gov^a allele of the CD109 gene, and the other for the Gov^b allele of the CD109 gene. The first probe (SEQ ID NO:13) overlaps the CD109 intron 18/exon 19 junction, binds to the Gov^a allele at nucleotides 935 - 974 of SEQ ID NO:5, and was tagged with the fluorescent dye 6-FAM. The second probe (SEQ ID NO:14), also overlapping the CD109 intron 18/exon 19 junction, binds to the Gov^b allele at the position corresponding to nucleotides 935 - 971 of SEQ ID NO:5, and was tagged with the fluorescent dye VIC. Genomic DNA was isolated from Gov phenotyped human peripheral blood leukocytes, and PCR/hybridization analysis was carried out using Tagman real-time PCR technology (Perkin Elmer). Genomic DNA was amplified using primers SEQ ID NO:11 and SEQ ID NO:12, with each reaction additionally containing 100 nM FAM-labelled Gova probe and 200 nM VIC-labelled Govb probe. Allelic discrimination, based on allele-specific fluorescence, was then determined using a post-PCR plate reader (Perkin Elmer). In all cases, PCR/fluorescence-based Gov genotyping correlated with the Gov phenotype, indicating that the A to C polymorphism at position 2108 (SEQ ID NO: 1) does indeed segregate with the Gov phenotype.

EXAMPLE 4: SSP Analysis of PCR Amplified Genomic DNA

To further confirm that the A to C polymorphism at position 2108 of the CD109 open reading frame (nucleotide 2108, SEQ ID NO:1; nucleotide 954, SEQ ID NO:5) segregates with the Gov phenotype, we also performed an alternative analysis involving SSCP analysis of PCR amplified genomic CD109 DNA. Two Gov allele-specific antisense oligonucleotides - SEQ ID NO:6 and SEQ ID NO:7 – differing by a single 3' nucleotide (and binding to sequence complementary to nucleotides 954 – 976 of SEQ ID NO:5, and of the Gov^b counterpart of SEQ

ID NO:5, respectively), were combined with a common sense primer – SEQ ID NO:8 binds within intron 18 and which corresponds to nucleotides 752 – 773 of SEQ ID NO:5, to amplify a 225 bp genomic DNA fragment containing the Gov polymorphic site from genomic DNA isolated from Gov^{aa}, Gov^{ab}, and Gov^{bb} individuals. In all cases, complete concordance between PCR-SSP analysis and Gov phenotyping was observed.

SEQUENCES:

SEQ ID NO: 1 consists of the entire 4335 nucleotide CD109 cDNA open reading frame encoding the Gov^a allele. The Gov^a allele comprises an A at nucleotide position 2108.

SEQ ID NO:2 consists of the entire 1445 aa protein sequence produced from CD109 Gov^a cDNA. The Gov^a allele comprises a Tyr at amino acid 703.

SEQ ID NO: 3 consists of the entire 4335 nucleotide CD109 cDNA open reading frame encoding the Gov^b allele. The Gov^b allele comprises a C at nucleotide position 2108.

SEQ ID NO: 4 consists of the entire 1445 aa protein sequence produced from the CD109 Gov^b cDNA. The Gov^b allele comprises a Ser at amino acid 703.

SEQ ID NO: 5 consists of the CD109 genomic DNA comprising CD109 exon 19 and the flanking introns, introns 18 and 19. The 118 nucleotide exon 19, comprising nucleotides 952 – 1069 of SEQ ID NO:5, corresponds to nucleotides 2106 – 2223 of SEQ ID NO: 1. The A to C Gov polymorphism of CD109 (corresponding to nucleotide 2108 of SEQ ID NO: 1) therefore corresponds to nucleotide 954 of SEQ ID NO:5. In the Gov^a allele, nucleotide 954 is A, while in the Gov^b allele nucleotide 954 is C. Thus, SEQ ID NO:5 corresponds to the Gov^a allele of CD109. Within SEQ ID NO:5, nucleotides 1 – 951 correspond to CD109 intron 18, while nucleotides 1070 – 2608 correspond to intron 19.

We note that nucleotides 2108 – 2112 of SEQ ID NO: 1, and the corresponding nucleotides 954 – 958 of SEQ ID NO:5, which consist of the sequence 5' ACAGG 3' (and which contains the Gov^a allele-specific polymorphic nucleotide at its 5' end), is not cleavable by the restriction endonuclease BstNI. However, in the corresponding Gov^b allele, the corresponding sequence – 5' CCAGG 3' – is cleavable by BstNI, and that the two Gov alleles can be discriminated on this basis. We note also that a group of restriction endonucleases - Bst2UI,

BstNI, BstOI, EcoRII, MaeIII, MspR91, MvaI, or ScrFI (or one of their isoschizomers) – is capable of differentiating between the Gov^a and Gov^b alleles on this basis.

SEQ ID NO:6 – **SEQ ID NO:14** comprise oligonucleotides for the PCR amplification of Gov polymorphism containing CD109 sequence from RNA, cDNA derived from RNA, or from genomic DNA, and for the Gov typing analyses of such amplified DNA fragments.

SEQ ID NO:6.

SEQ ID NO: 3, an antisense oligonucleotide specific for the Gov^a allele, binds to exon 19 sequence complementary to nucleotides 954 – 976 of SEQ ID NO:5. SEQ ID NO:6 and SEQ ID NO:7 (see below) differ by a single allele-specific 3' nucleotide

SEQ ID NO:7.

SEQ ID NO:7, an antisense oligonucleotide specific for the Gov^b allele, binds to exon 19 sequence complementary to nucleotides 954 – 976 of the Gov^b counterpart of SEQ ID NO:5. SEQ ID NO:6 (see above) and SEQ ID NO:7 differ by a single allele-specific 3' nucleotide.

SEQ ID NO:8.

SEQ ID NO:8 binds within intron 18, and corresponds to nucleotides 752 - 773 of SEQ ID NO:5.

SEQ ID NO:9.

SEQ ID NO:9 binds within intron 18 (nucleotides 875 – 892 SEQ ID NO:5).

SEQ ID NO:10.

SEQ ID NO:10 binds within intron 19 to the sequence complementary to nucleotides 1305 – 1322 of SEQ ID NO:5.

SEQ ID NO:11

SEQ ID NO:11 binds within intron 18 to nucleotides 902 – 928 of SEQ ID NO:5.

SEQ ID NO:12.

SEQ ID NO:12, binds within exon 19 to the sequence complementary to nucleotides 977 – 1006 of SEQ ID NO:5.

SEQ ID NO:13.

SEQ ID NO:13, specific for the Gov^a allele, overlaps the CD109 intron 18/exon 19 junction, and binds to the Gov^a allele at nucleotides 935 – 974 of SEQ ID NO:5.

SEQ ID NO:14.

SEQ ID NO:14, specific for the Gov^b allele, overlaps the CD109 intron 18/exon 19 junction, and binds to the Gov^b allele at the position corresponding to nucleotides 935 – 971 of SEQ ID NO:5.

SEQUENCE LISTING

<110> SCHUH, Andre and OUWEHAND, Willem

<120> Diagnosis and Treatment of Blood Disorders

<130> 3170/0005

<160> 30

<170> PatentIn version 3.0

<210> 1

<211> 4335

<212> DŅA

<213> HUMAN

<220>

<221> CDS

<222> (1)..(4335)

	0> 1 cag Gln	ggc														48	
	gcc Ala															96	
cca Pro	gly ggg	atc Ile 35	atc Ile	agg Arg	ccc Pro	gga Gly	gga Gly 40	aat Asn	gtg Val	act Thr	att Ile	999 Gly 45	gtg Val	gag Glu	ctt Leu	144	
	gaa Glu 50															192	
	gca Ala															240	

		Ser						cta Leu			288
								cgt Arg			336
								gag Glu 125			384
								aag Lys		caa Gln	432
								ttt Phe			480
					_	_		tca Ser	_		528
_		_		_	_		 _	att Ile			576
								tct Ser 205			624
								gtt Val			672
								tta Leu			720
								aag Lys			768
	${\tt Pro}$			Val		Leu		tta Leu			816
								aag Lys 285			. 864
								aat Asn			912
							Ser	tcc Ser			960
			Thr					ggt Gly			1008

aat Asn	gta Val	agc Ser	act Thr 340	aat Asn	gtg Val	ttc Phe	ttc Phe	aag Lys 345	caa Gln	cat His	gat Asp	tac Tyr	atc Ile 350	att Ile	gag Glu	1056
ttt Phe	ttt Phe	gat Asp 355	tat Tyr	act Thr	act Thr	gtc Val	ttg Leu 360	aag Lys	cca Pro	tct Ser	ctc Leu	aac Asn 365	ttc Phe	aca Thr	gcc Ala	1104
act Thr	gtg Val 370	aag Lys	gta Val	act Thr	cgt Arg	gct Ala 375	gat Asp	ggc Gly	aac Asn	caa Gln	ctg Leu 380	act Thr	ctt Leu	gaa Glu	gaa Glu	1152
aga Arg 385	aga Arg	aat Asn	aat Asn	gta Val	gtc Val 390	ata Ile	aca Thr	gtg Val	aca Thr	cag Gln 395	aga Arg	aac Asn	tat Tyr	act Thr	gag Glu 400	1200
tac Tyr	tgg Trp	agc Ser	gga Gly	tct Ser 405	aac Asn	agt Ser	gga Gly	aat Asn	cag Gln 410	aaa Lys	atg Met	gaa Glu	gct Ala	gtt Val 415	cag Gln	1248
aaa Lys	ata Ile	aat Asn	tat Tyr 420	act Thr	gtc Val	ccc Pro	caa Gln	agt Ser 425	gga Gly	act Thr	ttt Phe	aag Lys	att Ile 430	gaa Glu	ttc Phe	1296
cca Pro	atc Ile	ctg Leu 435	gag Glu	gat Asp	tcc Ser	agt Ser	gag Glu 440	cta Leu	cag Gln	ttg Leu	aag Lys	gcc Ala 445	tat Tyr	ttc Phe	ctt Leu	1344
ggt Gly	agt Ser 450	ааа L ув	agt Ser	agc Ser	atg Met	gca Ala 455	gtt Val	cat His	agt Ser	ctg Leu	ttt Phe 460	aag Lys	tct Ser	cct Pro	agt Ser	1392
aag Lys 465	aca Thr	tac Tyr	atc Ile	caa Gln	cta Leu 470	aaa Lys	aca Thr	aga Arg	gat Asp	gaa Glu 475	aat Asn	ata Ile	aag Lys	gtg Val	gga Gly 480	. 1440
tcg Ser	cct Pro	ttt Phe	gag Glu	ttg Leu 485	gtg Val	gtt Val	agt Ser	ggc Gly	aac Asn 490	aaa Lys	cga Arg	ttg Leu	aag Lys	gag Glu 495	tta Leu	1488
agc Ser	tat Tyr	atg Met	gta Val 500	gta Val	tcc Ser	agg Arg	gga Gly	cag Gln 505	ttg Leu	gtg Val	gct Ala	gta Val	gga Gly 510	aaa Lys	caa Gln	1536
aat Asn	tca Ser	aca Thr 515	atg Met	ttc Phe	tct Ser	tta Leu	aca Thr 520	cca Pro	gaa Glu	aat Asn	tct Ser	tgg Trp 525	act Thr	cca Pro	aaa Lys	1584
gcc Ala	tgt Cys 530	gta Val	att Ile	gtg Val	tat Tyr	tat Tyr 535	att Ile	gaa Glu	gat Asp	gat Asp	999 Gly 540	gaa Glu	att Ile	ata Ile	agt Ser	1632
gat Asp 545	gtt Val	cta Leu	aaa Lys	att Ile	cct Pro 550	gtt Val	cag Gln	ctt Leu	gtt Val	ttt Phe 555	aaa Lys	aat Asn	aag Lys	ata Ile	aag Lys 560	1680
cta Leu	tat Tyr	tgg Trp	agt Ser	aaa Lys 565	gtg Val	aaa Lys	gct Ala	gaa Glu	cca Pro 570	tct Ser	gag Glu	ааа Lyв	gtc Val	tct Ser 575	ctt Leu	1728
agg Arg	atc Ile	tct Ser	gtg Val 580	aca Thr	cag Gln	cct Pro	gac Asp	tcc Ser 585	ata Ile	gtt Val	Gly 888	att Ile	gta Val 590	gct Ala	gtt Val	1776

gac Asp	aaa Lys	agt Ser 595	gtg Val	aat Asn	ctg Leu	atg Met	aat Asn 600	gcc Ala	tct Ser	aat Asn	gat Asp	att Ile 605	aca Thr	atg Met	gaa Glu	1824
aat Asn	gtg Val 610	gtc Val	cat His	gag Glu	ttg Leu	gaa Glu 615	ctt Leu	tat Tyr	aac Asn	aca Thr	gga Gly 620	tat Tyr	tat Tyr	tta Leu	ggc	1872
			aat Asn													1920
			gca Ala													1968
			tat Tyr 660													2016
			cat His													2064
			gag Glu													2112
			gaa Glu													2160
gtg Val	gct Ala	act Thr	ggt Gly	ttt Phe 725	gtg Val	atc Ile	tct Ser	gag Glu	gac Asp 730	ctg Leu	ggt Gly	ctt Leu	gga Gly	cta Leu 735	aca Thr	2208
			gtg Val 740													2256
aat Asn	ctt Leu	ccc Pro 755	tac Tyr	tct Ser	gtt Val	atc Ile	aga Arg 760	ggt Gly	gaa Glu	gaa Glu	ttt Phe	gct Ala 765	ttg Leu	gaa Glu	ata Ile	2304
act Thr	ata Ile 770	ttc Phe	aat Asn	tat Tyr	ttg Leu	ааа L ys 775	gat Asp	gcc Ala	act Thr	gag Glu	gtt Val 780	aag Lys	gta Val	atc Ile	att Ile	2352
gag Glu 785	aaa Lys	agt Ser	gac Asp	aaa Lys	ttt Phe 790	gat Asp	att Ile	cta Leu	atg Met	act Thr 795	tca Ser	aat Asn	gaa Glu	ata Ile	aat Asn 800	2400
gcc Ala	aca Thr	ggc Gly	cac His	cag Gln 805	Gln	acc Thr	ctt Leu	ctg Leu	gtt Val 810	ccc Pro	agt Ser	gag Glu	Asp	999 Gly 815	gca Ala	2448
			ttt Phe 820													2496
			gct Ala													2544

	tta Leu 850															2592
	tta Leu															2640
	ttc Phe															2688
	act Thr													Leu		2736
	ttg Leu															2784
	gct Ala 930															2832
	aca Thr															2880
	cag Gln															2928
	gly ggg															2976
								Pro					À		ag aat ln Asn	3024
	tta Leu 1010	His			tac Tyr		Tr				y Hi		_	aaa t Lys s		3069
	ggt Gly 1025	Glu			gat Asp		ĞĨ							gag (31u I		3114
	ggt Gly 1040	Gly					Va				r Āl					3159
	tct Ser 1055	Let					Lу				o As					3204
caa Gln	gag Glu 1070	Ser	ato Ile	cat His	ttt Phe	ttg Leu 107	Ğl	g to u Se	r Gl	u Ph	c ag e Se . 10	r A	ga g rg (gga a Bly l	itt [le	3249
tca Ser	gac Asp 1085	Asn	tat Tyr	act Thr	cta Leu	gcc Ala 109	Le	t at u Il	a ac e Th	t ta r Ty	r Al	a t a L 95	tg t eu s	ca t Ser S	ca Ser	3294

	999 Gly 1100						Glu				atg Met 1110				3339
	gca Ala 1115						Met				gtg Val 1125	Ser			3384
	aaa Lys 1130	Leu	tct Ser	gac Asp	tcc Ser	tgg Trp 1135	Gln	cca Pro	cgc Arg	tcc Ser	ctg Leu 1140	gat Asp	att Ile	gaa Glu	3429
	gca Ala 1145														3474
	gag Glu 1160														3519
agc Ser	ttg Leu 1175	Gly	ggt Gly	ttt Phe	gca' Ala	tct Ser 1180	Thr	cag Gln	gat Asp	acc Thr	act Thr 1185	gtg Val	gct Ala	tta Leu	3564
	gct Ala 1190														3609
	atc Ile 1205					_			_			-			3654
_	ttt Phe 1220										ctt Leu 1230				3699
	ctt Leu 1235										att Ile 1245		gca Ala		3744
	ttt Phe 1250												aat Asn		3789
	gct Ala 1265														3834
gaa Glu	gcc Ala 1280	ttt Phe	gat Asp	tta Leu	gat Asp	gtt Val 1285	gct Ala	gta Val	aaa Lys	gaa Glu	aat Asn 1290	aaa Lys	gat Asp	gat . Asp	3879
	aat Asn 1295	His	gtg Val	gat Asp	ttg Leu	aat Asn 1300	gtg Val	tgt Cys	aca Thr	agc Ser	ttt Phe 1305	tcg Ser	ggc Gly	ccg Pro	3924
	agg Arg 1310	Ser													3969
	atg Met 1325														4014

aaa Lys	gtg Val 1340	gaa Glu	tat Tyr	gat Asp	cat His	gga Gly 1345	aaa Lys	ctc Leu	aac Asn	ctc Leu	tat Tyr 1350	tta Leu	gat Asp	tct Ser	4059
gta Val	aat Asn 1355	gaa Glu	acc Thr	cag Gln	ttt Phe	tgt Cys 1360	Val	aat Asn	att Ile	cct Pro	gct Ala 1365	gtg Val	aga Arg	aac Asn	4104
ttt Phe	aaa Lys 1370	Val	tca Ser	aat Asn	acc Thr	caa Gln 1375	Asp	gct Ala	tca Ser	gtg Val	tcc Ser 1380	ata Ile	gtg Val	gat Asp	4149
tac Tyr	tat Tyr 1385	gag Glu	cca Pro	agg Arg	aga Arg	cag Gln 1390	gcg Ala	gtg Val	aga Arg	agt Ser	tac Tyr 1395	aac Asn	tct Ser	gaa Glu	4194
	aag Lys 1400	ctg Leu	tcc Ser	tcc Ser	tgt Cys	gac Asp 1405	ctt Leu	tgc Cys	agt Ser	gat Asp	gtc Val 1410	cag Gln	ggc Gly	tgc Cys	4239
cgt Arg	cct Pro 1415	tgt Cys	gag Glu	gat Asp	gga Gly	gct Ala 1420	tca Ser	ggc Gly	tcc Ser	cat His	cat His 1425	cac His	tct Ser	tca Ser	4284
	att Ile 1430	ttt Phe	att Ile	ttc Phe	tgt Cys	ttc Phe 1435	aag Lys	ctt Leu	ctg Leu	tac Tyr	ttt Phe 1440	atg Met	gaa Glu	ctt Leu	4329
	ctg Leu 1445										•				4335

<210> 2

<211> 1445

<212> PRT

<213> HUMAN

<400> 2

Met Gln Gly Pro Pro Leu Leu Thr Ala Ala His Leu Leu Cys Val Cys
1 10 15

Thr Ala Ala Leu Ala Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala 20 25 30

Pro Gly Ile Ile Arg Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu 35 40 45

Leu Glu His Cys Pro Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys 50 55 60

Thr Ala Ser Asn Leu Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe 65 70 75 80

Glu Lys Gly Ser Phe Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn 85 90 95

- Ser Ala Asp Glu Ile Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp 100 105 110
- Glu Ile Leu Phe Ser Asn Ser Thr Arg Leu Ser Phe Glu Thr Lys Arg 115 120 125
- Ile Ser Val Phe Ile Gln Thr Asp Lys Ala Leu Tyr Lys Pro Lys Gln 130 135 140
- Glu Val Lys Phe Arg Ile Val Thr Leu Phe Ser Asp Phe Lys Pro Tyr 145 150 155 160
- Lys Thr Ser Leu Asn Ile Leu Ile Lys Asp Pro Lys Ser Asn Leu Ile 165 170 175
- Gln Gln Trp Leu Ser Gln Gln Ser Asp Leu Gly Val Ile Ser Lys Thr 180 185 190
- Phe Gln Leu Ser Ser His Pro Ile Leu Gly Asp Trp Ser Ile Gln Val 195 200 205
- Gln Val Asn Asp Gln Thr Tyr Tyr Gln Ser Phe Gln Val Ser Glu Tyr 210 215 220
- Met Asn Ser Lys His Leu Asn Gly Thr Ile Thr Ala Lys Tyr Thr Tyr 245 250 255
- Gly Lys Pro Val Lys Gly Asp Val Thr Leu Thr Phe Leu Pro Leu Ser 260 265 270
- Phe Trp Gly Lys Lys Asn Ile Thr Lys Thr Phe Lys Ile Asn Gly 275 280 285
- Ser Ala Asn Phe Ser Phe Asn Asp Glu Glu Met Lys Asn Val Met Asp 290 295 300
- Ser Ser Asn Gly Leu Ser Glu Tyr Leu Asp Leu Ser Ser Pro Gly Pro 305 310 315 320
- Val Glu Ile Leu Thr Thr Val Thr Glu Ser Val Thr Gly Ile Ser Arg 325 330 335

Asn Val Ser Thr Asn Val Phe Phe Lys Gln His Asp Tyr Ile Ile Glu 340 345 350

- Phe Phe Asp Tyr Thr Thr Val Leu Lys Pro Ser Leu Asn Phe Thr Ala 355 360 365
- Thr Val Lys Val Thr Arg Ala Asp Gly Asn Gln Leu Thr Leu Glu Glu 370 375 380
- Arg Arg Asn Asn Val Val Ile Thr Val Thr Gln Arg Asn Tyr Thr Glu 385 395 400
- Tyr Trp Ser Gly Ser Asn Ser Gly Asn Gln Lys Met Glu Ala Val Gln 405 410 415
- Lys Ile Asn Tyr Thr Val Pro Gln Ser Gly Thr Phe Lys Ile Glu Phe 420 425 430
- Pro Ile Leu Glu Asp Ser Ser Glu Leu Gln Leu Lys Ala Tyr Phe Leu 435 440 445
- Gly Ser Lys Ser Ser Met Ala Val His Ser Leu Phe Lys Ser Pro Ser 450 455 460
- Lys Thr Tyr Ile Gln Leu Lys Thr Arg Asp Glu Asn Ile Lys Val Gly 465 470 475 480
- Ser Pro Phe Glu Leu Val Val Ser Gly Asn Lys Arg Leu Lys Glu Leu 485 490 495
- Ser Tyr Met Val Val Ser Arg Gly Gln Leu Val Ala Val Gly Lys Gln 500 505 510
- Asn Ser Thr Met Phe Ser Leu Thr Pro Glu Asn Ser Trp Thr Pro Lys 515 520 525
- Ala Cys Val Ile Val Tyr Tyr Ile Glu Asp Asp Gly Glu Ile Ile Ser .530 535 540
- Asp Val Leu Lys Ile Pro Val Gln Leu Val Phe Lys Asn Lys Ile Lys 545 555 560
- Leu Tyr Trp Ser Lys Val Lys Ala Glu Pro Ser Glu Lys Val Ser Leu 565 570 575
- Arg Ile Ser Val Thr Gln Pro Asp Ser Ile Val Gly Ile Val Ala Val 580 585 590

Asp Lys Ser Val Asn Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu 595 600 605

- Asn Val Val His Glu Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly 610 615 620
- Met Phe Met Asn Ser Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val 625 630 635
- Leu Thr Asp Ala Asn Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp 645 650 655
- Asn Ala Glu Tyr Ala Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile 660 665 670
- Val Asp Ile His Asp Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys 675 680 685
- His Phe Pro Glu Thr Trp Ile Trp Leu Asp Thr Asn Met Gly Tyr Arg 690 695 700
- Ile Tyr Gln Glu Phe Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp
 705 710 715 720
- Val Ala Thr Gly Phe Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr 725 730 735
- Thr Thr Pro Val Glu Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu 740 745 750
- Asn Leu Pro Tyr Ser Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile . 755 760 765
- Thr Ile Phe Asn Tyr Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile 770 775 780
- Glu Lys Ser Asp Lys Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn 785 790 795 800
- Ala Thr Gly His Gln Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala 805 810 815
- Thr Val Leu Phe Pro Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile 820 825 830
- Thr Val Thr Ala Leu Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met 835 840 845

Ile Leu Val Lys Ala Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile 850 855 860

- Leu Leu Asp Leu Thr Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu 865 · 870 875 880
- Ser Phe Ser Phe Pro Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln 885 890 895
- Ile Thr Ala Ile Gly Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala 900 905 910
- Ser Leu Ile Arg Met Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn 915 920 925
- Phe Ala Pro Asn Ile Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Gln 930 935 940
- Leu Thr Asp Asn Leu Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly 945 950 955 960
- Tyr Gln Arg Glu Leu Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala 965 970 975
- Phe Gly Asn Tyr Asp Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val 980 985 990
- Leu Arg Cys Phe Leu Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn 995 1000 1005
- Val Leu His Arg Thr Tyr Thr Trp Leu Lys Gly His Gln Lys Ser 1010 1015 1020
- Asn Gly Glu Phe Trp Asp Pro Gly Arg Val Ile His Ser Glu Leu 1025 1030 1035
- Gln Gly Gly Asn Lys Ser Pro Val Thr Leu Thr Ala Tyr Ile Val 1040 1045 1050
- Thr Ser Leu Leu Gly Tyr Arg Lys Tyr Gln Pro Asn Ile Asp Val 1055 1060 1065
- Gln Glu Ser Ile His Phe Leu Glu Ser Glu Phe Ser Arg Gly Ile 1070 1075 1080
- Ser Asp Asn Tyr Thr Leu Ala Leu Ile Thr Tyr Ala Leu Ser Ser 1085 1090 1095

Val Gly Ser Pro Lys Ala Lys Glu Ala Leu Asn Met Leu Thr Trp . 1105 Arg Ala Glu Gln Glu Gly Gly Met Gln Phe Trp Val Ser Ser Glu 1120 1125 1115 Ser Lys Leu Ser Asp Ser Trp Gln Pro Arg Ser Leu Asp Ile Glu 1130 1135 · Val Ala Ala Tyr Ala Leu Leu Ser His Phe Leu Gln Phe Gln Thr 1150 Ser Glu Gly Ile Pro Ile Met Arg Trp Leu Ser Arg Gln Arg Asn 1165 1160 Ser Leu Gly Gly Phe Ala Ser Thr Gln Asp Thr Thr Val Ala Leu Lys Ala Leu Ser Glu Phe Ala Ala Leu Met Asn Thr Glu Arg Thr 1195 1190 Asn Ile Gln Val Thr Val Thr Gly Pro Ser Ser Pro Ser Pro Val 1205 Lys Phe Leu Ile Asp Thr His Asn Arg Leu Leu Leu Gln Thr Ala 1220 · 1225 Glu Leu Ala Val Val Gln Pro Met Ala Val Asn Ile Ser Ala Asn 1240 1245 Gly Phe Gly Phe Ala Ile Cys Gln Leu Asn Val Val Tyr Asn Val 1255 Lys Ala Ser Gly Ser Ser Arg Arg Arg Arg Ser Ile Gln Asn Gln 1270 Glu Ala Phe Asp Leu Asp Val Ala Val Lys Glu Asn Lys Asp Asp 1285 1280 Leu Asn His Val Asp Leu Asn Val Cys Thr Ser Phe Ser Gly Pro 1295 1300 1305 Gly Arg Ser Gly Met Ala Leu Met Glu Val Asn Leu Leu Ser Gly 1310 Phe Met Val Pro Ser Glu Ala Ile Ser Leu Ser Glu Thr Val Lys 1330 1325

Lys Val Glu Tyr Asp His Gly Lys Leu Asn Leu Tyr Leu Asp Ser 1340 : 1345 : 1350

- Val Asn Glu Thr Gln Phe Cys Val Asn Ile Pro Ala Val Arg Asn 1355 1360 1365
- Phe Lys Val Ser Asn Thr Gln Asp Ala Ser Val Ser Ile Val Asp 1370 1375 1380
- Tyr Tyr Glu Pro Arg Arg Gln Ala Val Arg Ser Tyr Asn Ser Glu 1385 1390 1395
- Val Lys Leu Ser Ser Cys Asp Leu Cys Ser Asp Val Gln Gly Cys 1400 1405 1410
- Arg Pro Cys Glu Asp Gly Ala Ser Gly Ser His His Ser Ser 1415 1420 1425
- Val Ile Phe Ile Phe Cys Phe Lys Leu Leu Tyr Phe Met Glu Leu 1430 1440
- Trp Leu 1445
- <210> 3
- <211> 4335
- <212> DNA
- <213> HUMAN
- <220>
- <221> CDS
- <222> (1)..(4335)
- <220>
- <221> CDS
- <222> (1)..(4335)

48

acc g Thr A	la la	gcg Ala	ctg Leu 20	gcc Ala	gtg Val	gct Ala	ccc Pro	999 Gly 25	cct Pro	cgg Arg	ttt Phe	ctg Leu	gtg Val 30	aca Thr	gcc Ala		96
cca g Pro G	31 y 399	atc Ile 35	atc Ile	agg Arg	ccc Pro	gga Gly	gga Gly 40	aat Asn	gtg Val	act Thr	att Ile	999 Gly 45	gtg Val	gag Glu	ctt Leu		144
ctg g Leu G 5	gaa 31u 50	cac Kis	tgc Cys	cct Pro	tca Ser	cag Gln 55	gtg Val	act Thr	gtg Val	aag Lys	gcg Ala 60	gag Glu	ctg Leu	ctc Leu	aag Lys		192
aca g Thr A 65	gca Ma	tca Ser	aac Asn	ctc Leu	act Thr 70	gtc Val	tct Ser	gtc Val	ctg Leu	gaa Glu 75	gca Ala	gaa Glu	gga Gly	gtc Val	ttt Phe 80		240
gaa a Glu L	aaa Lys	ggc Gly	tct Ser	ttt Phe 85	aag Lys	aca Thr	ctt Leu	act Thr	ctt Leu 90	cca Pro	tca Ser	cta Leu	cct Pro	ctg Leu 95	aac Asn		288
agt g Ser A	gca Ala	gat Asp	gag Glu 100	att Ile	tat Tyr	gag Glu	cta Leu	cgt Arg 105	gta Val	acc Thr	gga Gly	cgt Arg	acc Thr 110	cag Gln	gat Asp		336
gag a Glu I	att [le	tta Leu 115	ttc Phe	tct Ser	aat Asn	agt Ser	acc Thr 120	cgc Arg	tta Leu	tca Ser	ttt Phe	gag Glu 125	acc Thr	aag Lys	aga Arg		384
ata t Ile S 1																	432
gaa g Glu V 145	gtg /al	aag Lys	ttt Phe	cgc Arg	att Ile 150	gtt Val	aca Thr	ctc Leu	ttc Phe	tca Ser 155	gat Asp	ttt Phe	aag Lys	cct Pro	tac Tyr 160		480
aaa a Lys T																	528
caa c Gln G	eag 31n	tgg Trp	ttg Leu 180	tca Ser	caa Gln	caa Gln	agt Ser	gat Asp 185	ctt Leu	gga Gly	gtc Val	att Ile	tcc Ser 190	aaa Lys	act Thr		576
ttt c Phe G																	624 ·
caa g Gln V 2	gtg Val 210	aat Asn	gac Asp	cag Gln	aca Thr	tat Tyr 215	tat Tyr	caa Gln	tca Ser	ttt Phe	cag Gln 220	gtt Val	tca Ser	gaa Glu	tat Tyr		672
gta t Val L 225	tta Leu	cca Pro	aaa Lys	ttt Phe	gaa Glu 230	gtg Val	act Thr	ttg Leu	cag Gln	aca Thr 235	cca Pro	tta Leu	tat Tyr	tgt Cys	tct Ser 240		720
atg a Met A	aat Asn	tct Ser	aag Lys	cat His 245	tta Leu	aat Asn	ggt Gly	acc Thr	atc Ile 250	acg Thr	gca Ala	aag Lys	tat Tyr	aca Thr 255	tat Tyr	,	768
ggg a Gly L	aag Lys	cca Pro	gtg Val 260	aaa Lys	gga Gly	gac Asp	gta Val	acg Thr 265	ctt Leu	aca Thr	ttt Phe	tta Leu	cct Pro 270	tta Leu	tcc Ser		816

ttt Phe	tgg Trp	gga Gly 275	aag Lys	aag Lys	aaa Lys	aat Asn	att Ile 280	aca Thr	aaa Lys	aca Thr	ttt Phe	aag Lys 285	ata Ile	aat Asn	gga Gly	864
tct Ser	gca Ala 290	aac Asn	ttc Phe	tct Ser	ttt Phe	aat Asn 295	gat Asp	gaa Glu	gag Glu	atg Met	aaa Lys 300	aat Asn	gta Val	atg Met	gat Asp	912
								ctg Leu								960
gta Val	gaa Glu	att Ile	tta Leu	acc Thr 325	aca Thr	gtg Val	aca Thr	gaa Glu	tca Ser 330	gtt Val	aca Thr	ggt Gly	att Ile	tca Ser 335	aga Arg	1008
aat Asn	gta Val	agc Ser	act Thr 340	aat Asn	gtg Val	ttc Phe	ttc Phe	aag Lys 345	caa Gln	cat His	gat Asp	tac Tyr	atc Ile 350	att Ile	gag Glu	1056
ttt Phe	ttt Phe	gat Asp 355	Tyr	act Thr	act Thr	gtc Val	ttg Leu 360	aag Lys	cca Pro	tct Ser	ctc Leu	aac Asn 365	ttc Phe	aca Thr	gcc Ala	1104
act Thr	gtg Val 370	aag Lys	gta Val	act Thr	cgt Arg	gct Ala 375	gat Asp	Gly	aac Asn	caa Gln	ctg Leu 380	act Thr	ctt Leu	gaa Glu	gaa Glu	1152
								gtg Val								1200
tac Tyr	tgg Trp	agc Ser	gga Gly	tct Ser 405	aac Asn	agt Ser	gga Gly	aat Asn	cag Gln 410	aaa Lys	atg Met	gaa Glu	gct Ala	gtt Val 415	cag Gln	1248
aaa Lys	ata Ile	aat Asn	tat Tyr 420	act Thr	gtc Val	ccc Pro	caa Gln	agt Ser 425	gga Gly	act Thr	ttt Phe	aag Lys	att Ile 430	gaa Glu	ttc Phe	1296
cca Pro	atc Ile	ctg Leu 435	gag Glu	gat Asp	tcc Ser	agt Ser	gag Glu 440	cta Leu	cag Gln	ttg Leu	aag Lys	gcc Ala 445	tat Tyr	ttc Phe	ctt Leu	1344
ggt Gly	agt Ser 450	aaa Lys	agt Ser	agc Ser	atg Met	gca Ala 455	gtt Val	cat His	Ser	ctg Leu	ttt Phe 460	aag Lys	tct Ser	cct Pro	agt Ser	1392
aag Lys 465	aca Thr	tac Tyr	atc Ile	caa Gln	cta Leu 470	aaa Lys	aca Thr	aga Arg	gat Asp	gaa Glu 475	aat Asn	ata Ile	aag Lys	gtg Val	gga Gly 480	1440
tcg Ser	cct Pro	ttt Phe	gag Glu	ttg Leu 485	gtg Val	gtt Val	agt Ser	ggc Gly	aac Asn 490	aaa Lys	cga Arg	ttg Leu	aag Lys	gág Glu 495	tta Leu	1488
agc Ser	tat Tyr	Met	gta Val ·500	gta Val	tcc Ser	agg Arg	gga Gly	cag Gln 505	ttg Leu	gtg Val	gct Ala	gta Val	gga Gly 510	aaa Lys	caa Gln	1536
aat Asn	tca Ser	aca Thr 515	atg Met	ttc Phe	tct Ser	tta Leu	aca Thr 520	cca Pro	gaa Glu	aat Asn	tct Ser	tgg Trp 525	act Thr	cca Pro	aaa Lys	1584

gcc tgt gta att Ala Cys Val Ile 530	gtg tat tat Val Tyr Tyr 535	att gaa gat Ile Glu Asp	gat ggg gaa at Asp Gly Glu Il 540	t ata agt 1632 e Ile Ser
gat gtt cta aaa Asp Val Leu Lys 545				
cta tat tgg agt Leu Tyr Trp Ser				
agg atc tct gtg Arg Ile Ser Val 580	aca cag cet Thr Gln Pro	gac tcc ata Asp Ser Ile 585	gtt ggg att gt Val Gly Ile Va 59	l Ala Val
gac aaa agt gtg Asp Lys Ser Val 595	aat ctg atg Asn Leu Met	aat gcc tct Asn Ala Ser 600	aat gat att ac Asn Asp Ile Th 605	a atg gaa 1824 r Met Glu
aat gtg gtc cat Asn Val Val His 610				
atg ttc atg aat Met Phe Met Asn 625				
ttg aca gat gca Leu Thr Asp Ala	aac ctc acg Asn Leu Thr 645	aag gat tat Lys Asp Tyr 650	att gat ggt gt Ile Asp Gly Va	t tat gac 1968 l Tyr Asp 655
aat gca gaa tat Asn Ala Glu Tyr 660				y His Ile
gta gat att cat Val Asp Ile His 675				
cat ttt cca gag His Phe Pro Glu 690	act tgg att Thr Trp Ile 695	tgg cta gac Trp Leu Asp	acc aac atg gg Thr Asn Met Gl 700	t toe agg 2112 y Ser Arg
att tac caa gaa Ile Tyr Gln Glu 705	ttt gaa gta Phe Glu Val 710	act gta cct Thr Val Pro	gat tct atc ac Asp Ser Ile Th 715	t tct tgg 2160 r Ser Trp 720
gtg gct act ggt Val Ala Thr Gly				
act act cca gtg Thr Thr Pro Val 740				e Phe Leu
aat ctt ccc tac Asn Leu Pro Tyr 755				
act ata ttc aat Thr Ile Phe Asn 770	tat ttg aaa Tyr Leu Lys 775	gat gcc act Asp Ala Thr	gag gtt aag gt Glu Val Lys Va 780	a atc att 2352 l Ile Ile

gag aaa agt gac aaa ttt gat att cta atg act tca aat gaa ata aat Glu Lys Ser Asp Lys Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn 785 790 795 800	2400
gcc aca ggc cac cag cag acc ctt ctg gtt ccc agt gag gat ggg gca Ala Thr Gly His Gln Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala 805 810 815	2448
act gtt ctt ttt ccc atc agg cca aca cat ctg gga gaa att cct atc Thr Val Leu Phe Pro Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile 820 825 830	2496
aca gtc aca gct ctt tca ccc act gct tct gat gct gtc acc cag atg Thr Val Thr Ala Leu Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met 835 840 845	2544
att tta gta aag gct gaa gga ata gaa aaa tca tat tca caa tcc atc Ile Leu Val Lys Ala Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile 850 855 860	2592
tta tta gac ttg act gac aat agg cta cag agt acc ctg aaa act ttg Leu Leu Asp Leu Thr Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu 865 870 875 880	2640
agt ttc tca ttt cct cct aat aca gtg act ggc agt gaa aga gtt cag Ser Phe Ser Phe Pro Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln 885 890 895	2688
atc act gca att gga gat gtt ctt ggt cct tcc atc aat ggc tta gcc Ile Thr Ala Ile Gly Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala 900 910	2736
Ser Leu Ile Arg Met Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn 915 920 925	2784
ttt gct cca aat att tac att ttg gat tat ctg act aaa aag aaa caa Phe Ala Pro Asn Ile Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Gln 930 935 940	2832
ctg aca gat aat ttg aaa gaa aaa gct ctt tca ttt atg agg caa ggt Leu Thr Asp Asn Leu Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly 945 950 955 960	2880
tac cag aga gaa ctt ctc tat cag agg gaa gat ggc tct ttc agt gct Tyr Gln Arg Glu Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala 965 970 975	2928
ttt ggg aat tat gac cct tct ggg agc act tgg ttg tca gct ttt gtt Phe Gly Asn Tyr Asp Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val 980 985 990	2976
tta aga tgt ttc ctt gaa gcc gat cct tac ata gat att gat cag aat Leu Arg Cys Phe Leu Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn 995 1000 1005	3024
gtg tta cac aga aca tac act tgg ctt aaa gga cat cag aaa tcc Val Leu His Arg Thr Tyr Thr Trp Leu Lys Gly His Gln Lys Ser 1010 1015 1020	3069
aac ggt gaa ttt tgg gat cca gga aga gtg att cat agt gag ctt Asn Gly Glu Phe Trp Asp Pro Gly Arg Val Ile His Ser Glu Leu 1025 1030 1035	3114

Gln						cca Pro 1045									3159
Thr	tct Ser 1055	ctc Leu	ctg Leu	gga Gly	tat Tyr	aga Arg 1060	ГЛЗ	tat Tyr	cag Gln	cct Pro	aac Asn 1065	att Ile	gat Asp		3204
Gln						ttg Leu 1075									324J
Ser						gcc Ala 1090									3294
Val	999 Gly 1100					aag Lys 1105									3339
Arg						ggc Gly 1120									3384
Ser :						tgg Trp 1135									3429
Val 2						ctc Leu 1150	Ser								3474
Ser						atg Met 1165									3519
Ser 1						tct Ser 1180									3564
Lys i	gct Ala 1190	ctg Leu	tct Ser	gaa Glu	ttt Phe	gca Ala 1195	gcc Ala	cta Leu	atg Met	aat Asn	aca Thr 1200	gaa Glu	agg Arg	aca Thr	3609
Asn :	atc Ile 1205	caa Gln	gtg Val	acc Thr	gtg Val	acg Thr 1210	Gly 999	cct Pro	agc Ser	tca Ser	cca Pro 1215	agt Ser	cct Pro	gta Val	3654
Lys						cac His 1225									3699
Glu I	ctt Leu 1235	gct Ala	gtg Val	gta Val	cag Gln	cca Pro 1240	Met	gca Ala	gtt Val	aat Asn	att Ile 1245	tcc Ser	gca Ala	aat Asn	3744
Gly 1						tgt Cys 1255									3789
Lys i	gct Ala 1265	tct Ser	gly ggg	tct Ser	tct Ser	aga Arg 1270	aga Arg	cga Arg	aga Arg	tct Ser	atc Ile 1275	caa Gln	aat Asn	caa Gln	3834

gaa Glu	gcc Ala 1280	ttt Phe	gat Asp	tta Leu	gat Asp	gtt Val 1285	gct Ala	gta Val	aaa Lys	gaa Glu	aat Asn 1290	aaa Lys	gat Asp	gat Asp	3879
	aat Asn 1295										ttt Phe 1305	_	ggc	_	3924
ggt Gly	agg Arg 1310	agt Ser	ggc	atg Met	gct Ala	ctt Leu 1315	atg Met	gaa Glu	gtt Val	aac Asn	cta Leu 1320	tta Leu	agt Ser	ggc Gly	3969
											gag Glu 1335		gtg Val	_	4014
							Lys				tat Tyr 1350				4059
gta Val	aat Asn 1355	gaa Glu	acc Thr	cag Gln	ttt Phe	tgt Cys 1360	gtt Val	aat Asn	att Ile	cct Pro	gct Ala 1365	gtg Val	aga Arg	aac Asn	4104
ttt Phe	aaa Lys 1370	gtt Val	tca Ser	aat Asn	acc Thr	caa Gln 1375	gat Asp	gct Ala	tca Ser	gtg Val	tcc Ser 1380	ata Ile	gtg Val		4149
Tyr	Tyr 1385	Glu	Prò	Arg	Arg	Gln 1390	Ala	Val	Arg	Ser	tac Tyr 1395	Asn	Ser	Glu	4194
Val	Lys 1400	Leu	Ser	Ser	Сув	Asp 1405	Leu	Сув	Ser	Asp	gtc Val 1410	Gln	Gly	Cys	4239
Arg	Pro 1415	Сув	Glu	Āsp	Gly	Ala 1420	Ser	Gly	Ser	His	cat His 1425	His	Ser	Ser	4284
Val	Ile 1430										ttt Phe 1440				4329
tgg Trp	_														4335

<210> 4

<211> 1445

<212> PRT

<213> HUMAN

<400> 4

Met Gln Gly Pro Pro Leu Leu Thr Ala Ala His Leu Leu Cys Val Cys 1 5 10 15

Thr Ala Ala Leu Ala Val Ala Pro Gly Pro Arg Phe Leu Val Thr Ala 20 25 30

- Pro Gly Ile Ile Arg Pro Gly Gly Asn Val Thr Ile Gly Val Glu Leu 35 40 45
- Leu Glu His Cys Pro Ser Gln Val Thr Val Lys Ala Glu Leu Leu Lys 50 55 60
- Thr Ala Ser Asn Leu Thr Val Ser Val Leu Glu Ala Glu Gly Val Phe 65 70 75 80
- Glu Lys Gly Ser Phe Lys Thr Leu Thr Leu Pro Ser Leu Pro Leu Asn 85 90 95
- Ser Ala Asp Glu Ile Tyr Glu Leu Arg Val Thr Gly Arg Thr Gln Asp 100 105 110
- Glu Ile Leu Phe Ser Asn Ser Thr Arg Leu Ser Phe Glu Thr Lys Arg 115 120 125
- Ile Ser Val Phe Ile Gln Thr Asp Lys Ala Leu Tyr Lys Pro Lys Gln 130 135 140
- Glu Val Lys Phe Arg Ile Val Thr Leu Phe Ser Asp Phe Lys Pro Tyr 145 150 155 160
- Lys Thr Ser Leu Asn Ile Leu Ile Lys Asp Pro Lys Ser Asn Leu Ile 165 170 175
- Gln Gln Trp Leu Ser Gln Gln Ser Asp Leu Gly Val Ile Ser Lys Thr 180 185 190
- Phe Gln Leu Ser Ser His Pro Ile Leu Gly Asp Trp Ser Ile Gln Val 195 200 205
- Gln Val Asn Asp Gln Thr Tyr Tyr Gln Ser Phe Gln Val Ser Glu Tyr 210 215 220
- Val Leu Pro Lys Phe Glu Val Thr Leu Gln Thr Pro Leu Tyr Cys Ser 225 230 235 240
- Met Asn Ser Lys His Leu Asn Gly Thr Ile Thr Ala Lys Tyr Thr Tyr 245 250 255
- Gly Lys Pro Val Lys Gly Asp Val Thr Leu Thr Phe Leu Pro Leu Ser 260 265 270

Phe Trp Gly Lys Lys Lys Asn Ile Thr Lys Thr Phe Lys Ile Asn Gly 275 280 285

- Ser Ala Asn Phe Ser Phe Asn Asp Glu Glu Met Lys Asn Val Met Asp 290 295 300
- Ser Ser Asn Gly Leu Ser Glu Tyr Leu Asp Leu Ser Ser Pro Gly Pro 305 310 315
- Val Glu Ile Leu Thr Thr Val Thr Glu Ser Val Thr Gly Ile Ser Arg 325 330 335
- Asn Val Ser Thr Asn Val Phe Phe Lys Gln His Asp Tyr Ile Ile Glu 340 345 350
- Phe Phe Asp Tyr Thr Thr Val Leu Lys Pro Ser Leu Asn Phe Thr Ala 355 360 365
- Thr Val Lys Val Thr Arg Ala Asp Gly Asn Gln Leu Thr Leu Glu Glu 370 375 380
- Arg Arg Asn Asn Val Val Ile Thr Val Thr Gln Arg Asn Tyr Thr Glu 385 390 395 400
- Tyr Trp Ser Gly Ser Asn Ser Gly Asn Gln Lys Met Glu Ala Val Gln
 405 410 415
- Lys Ile Asn Tyr Thr Val Pro Gln Ser Gly Thr Phe Lys Ile Glu Phe 420 425 430
- Pro Ile Leu Glu Asp Ser Ser Glu Leu Gln Leu Lys Ala Tyr Phe Leu 435 440 445.
- Gly Ser Lys Ser Ser Met Ala Val His Ser Leu Phe Lys Ser Pro Ser 450 460
- Lys Thr Tyr Ile Gln Leu Lys Thr Arg Asp Glu Asn Ile Lys Val Gly
 465 470 475 480
- Ser Pro Phe Glu Leu Val Val Ser Gly Asn Lys Arg Leu Lys Glu Leu 485 490 495
- Ser Tyr Met Val Val Ser Arg Gly Gln Leu Val Ala Val Gly Lys Gln 500 505 510
- Asn Ser Thr Met Phe Ser Leu Thr Pro Glu Asn Ser Trp Thr Pro Lys 515 520 525

Ala Cys Val Ile Val Tyr Tyr Ile Glu Asp Asp Gly Glu Ile Ile Ser 530 535 540

- Asp Val Leu Lys Ile Pro Val Gln Leu Val Phe Lys Asn Lys Ile Lys 545 550 555
- Leu Tyr Trp Ser Lys Val Lys Ala Glu Pro Ser Glu Lys Val Ser Leu 565 570 575
- Arg Ile Ser Val Thr Gln Pro Asp Ser Ile Val Gly Ile Val Ala Val 580 585 590
- Asp Lys Ser Val Asn Leu Met Asn Ala Ser Asn Asp Ile Thr Met Glu 595 600 605
- Asn Val Val His Glu Leu Glu Leu Tyr Asn Thr Gly Tyr Tyr Leu Gly 610 615 620
- Met Phe Met Asn Ser Phe Ala Val Phe Gln Glu Cys Gly Leu Trp Val 625 630 635 640
- Leu Thr Asp Ala Asn Leu Thr Lys Asp Tyr Ile Asp Gly Val Tyr Asp 645 650 655
- Asn Ala Glu Tyr Ala Glu Arg Phe Met Glu Glu Asn Glu Gly His Ile 660 665 670
- Val Asp Ile His Asp Phe Ser Leu Gly Ser Ser Pro His Val Arg Lys 675 680 685
- His Phe Pro Glu Thr Trp Ile Trp Leu Asp Thr Asn Met Gly Ser Arg 690 695 700
- Ile Tyr Gln Glu Phe Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp 705 710 715 720
- Val Ala Thr Gly Phe Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr 725 730 735
- Thr Thr Pro Val Glu Leu Gln Ala Phe Gln Pro Phe Phe Ile Phe Leu 740 745 750
- Asn Leu Pro Tyr Ser Val Ile Arg Gly Glu Glu Phe Ala Leu Glu Ile 755 760 765
- Thr Ile Phe Asn Tyr Leu Lys Asp Ala Thr Glu Val Lys Val Ile Ile 770 780

Glu Lys Ser Asp Lys Phe Asp Ile Leu Met Thr Ser Asn Glu Ile Asn 785 790 795 800

- Ala Thr Gly His Gln Gln Thr Leu Leu Val Pro Ser Glu Asp Gly Ala 805 810 815
- Thr Val Leu Phe Pro Ile Arg Pro Thr His Leu Gly Glu Ile Pro Ile 820 825 830
- Thr Val Thr Ala Leu Ser Pro Thr Ala Ser Asp Ala Val Thr Gln Met 835 840 845
- Ile Leu Val Lys Ala Glu Gly Ile Glu Lys Ser Tyr Ser Gln Ser Ile 850 855 860
- Leu Leu Asp Leu Thr Asp Asn Arg Leu Gln Ser Thr Leu Lys Thr Leu 865 870 875 880
- Ser Phe Ser Phe Pro Pro Asn Thr Val Thr Gly Ser Glu Arg Val Gln 885 890 895
- Ile Thr Ala Ile Gly Asp Val Leu Gly Pro Ser Ile Asn Gly Leu Ala 900 905 910
- Ser Leu Ile Arg Met Pro Tyr Gly Cys Gly Glu Gln Asn Met Ile Asn 915 920 925
- Phe Ala Pro Asn Ile Tyr Ile Leu Asp Tyr Leu Thr Lys Lys Gln 930 940
- Leu Thr Asp Asn Leu Lys Glu Lys Ala Leu Ser Phe Met Arg Gln Gly 945 950 955 960
- Tyr Gln Arg Glu Leu Leu Tyr Gln Arg Glu Asp Gly Ser Phe Ser Ala 965 970 975
- Phe Gly Asn Tyr Asp Pro Ser Gly Ser Thr Trp Leu Ser Ala Phe Val 980 985 990
- Leu Arg Cys Phe Leu Glu Ala Asp Pro Tyr Ile Asp Ile Asp Gln Asn 995 1000 1005
- Val Leu His Arg Thr Tyr Thr Trp Leu Lys Gly His Gln Lys Ser 1010 1015
- Asn Gly Glu Phe Trp Asp Pro Gly Arg Val Ile His Ser Glu Leu 1025 1030 1035

									•							
(3ln	Gly 1040	_	Asn	Lys	Ser	Pro 1045		Thr	Leu	Thr	Ala 1050		Ile	Val	•
i	Thr	Ser 1055	Leu	Leu	Gly	Tyr	Arg 1060		Tyr	Gln	Pro	Asn 1065		Asp	Val	
(3ln	Glu 1070		Ile	His	Phe	Leu 1075		Ser	Glu	Phe	Ser 1080		Gly	Ile	
5	Ser	Asp 1085		Tyr	Thr	Leu	Ala 1090		Ile	Thr	Tyr	Ala 1095	Leu	Ser	Ser	
1	V al	Gly 1100	Ser	Pro	Lys	Ala	Lys 1105		Ala	Leu	Asn	Met . 1110	Leu	Thr	Trp	
1	Arg	Ala 1115	Glu	Gln	Glu	Gly	Gly 1120	Met	Gln	Phe	Trp	Val 1125	Ser	Ser	Glu	
8	Ser	Lys 1130	Leu	Ser	qaA		Trp 1135	Gln	Pro	Arg	Ser	Leu 1140	Asp	Ile	Glu	
7	Val	Ala 1145	Ala	Tyr	Ala	Leu	Leu 1150	Ser	His	Phe	Leu	Gln 1155	Phe	Gln	Thr	
5	Ser	Glu 1160	Gly	Ile	Pro	Ile	Met 1165	Arg	Trp	Leu	Ser	Arg 1170	Gln	Arg	Asn	
5	Ser	Leu 1175	Gly _.	Gly	Phe	Ala	Ser 1180	Thr	Gln	qaA	Thr	Thr 1185		Ala	Leu	
1	ρλâ	Ala 1190	Leu	Ser	Glu	Phe	Ala 1195	Ala	Leu	Met	Asn	Thr 1200	Glu	Arg	Thr .	
I		Ile 1205		Val	Thr		Thr 1210							Pro	Val	
1	ъув	Phe 1220	Leu	Ile	Asp	Thr	His 1225	Asn	Arg	Leu	Leu	Leu 1230	Gln	Thr	Ala	
C	3lu	Leu 1235	Ala	Val	Val	Gln	Pro 1240	Met	Ala		Asn	Ile 1245	Ser	Ala	Asn	
C	∃ly	Phe 1250	Gly	Phe	Ala	Ile	Cys 1255	Gln	Leu	Asn	Val	V al 1260	Tyr	Asn	Val	
I	у́s	Ala 1265	Ser	Gly	Ser	Ser	Arg 1270	Arg	Arg	Arg	Ser	Ile 1275	Gln	Asn	Gln	

Glu Ala Phe Asp Leu Asp Val Ala Val Lys Glu Asn Lys Asp Asp 1280 1285 1290

- Leu Asn His Val Asp Leu Asn Val Cys Thr Ser Phe Ser Gly Pro 1295 1300
- Gly Arg Ser Gly Met Ala Leu Met Glu Val Asn Leu Leu Ser Gly 1310 1315 1320
- Phe Met Val Pro Ser Glu Ala Ile Ser Leu Ser Glu Thr Val Lys 1325 1330 1335
- Lys Val Glu Tyr Asp His Gly Lys Leu Asn Leu Tyr Leu Asp Ser 1340 1345 1350
- Val Asn Glu Thr Gln Phe Cys Val Asn Ile Pro Ala Val Arg Asn 1355 1360 1365
- Phe Lys Val Ser Asn Thr Gln Asp Ala Ser Val Ser Ile Val Asp 1370 1375 1380
- Tyr Tyr Glu Pro Arg Arg Gln Ala Val Arg Ser Tyr Asn Ser Glu 1385 1390 1395
- Val Lys Leu Ser Ser Cys Asp Leu Cys Ser Asp Val Gln Gly Cys 1400 1410
- Arg Pro Cys Glu Asp Gly Ala Ser Gly Ser His His Ser Ser 1415 1420 1425
- Val Ile Phe Ile Phe Cys Phe Lys Leu Leu Tyr Phe Met Glu Leu 1430 1440

Trp Leu 1445

<210> 5

<211> 2608

<212> DNA

<213> HUMAN

<220>

<221> Intron

<222> (1)..(951)

<220>

<221> exon

<222> (952)..(1069)

<220>

<221> Intron

<222> (1070)..(2608)

<400> 5 60 gtaaaaattt ataaagttot ttgcccatac atattttgtt tagtgtttgt tttaaataag ctttgcccgc tttctaatgt ttaagtacaa acatagtgta actaagaact aagtagacca 120 aaaggatttt ttaggaaatg atatttattg aatctaaata cagtttttga taaagccaca 180 cataaattat ggcaggaagg tctcatcaat gagaagatag gcctttttt tttttttaa 240 ctgaagggtg attttgactt ccttgaagtc tcatgattct tgttgaagaa aaattgctgg 300 qaqtacattt gttgtcacag gatgggaagc actcatgatt acctcctgtg acccctggca 360 gtgctgctaa ctgaaccctg ctcctcacaa agcattccca ggagtcacag ggagaagggg 420 catqqqtqqt qqaaaqaatt caqcttggct gataaacccc gtaccacctg gcctgataat 480 540 tgagcaggta aatcatgaaa tccacatagt attttatagt cagctgttta aagatacttg agttaacaca tgagtgaaat ctcaaggaaa caaataacag cattgacagg gatacagaga 600 660 ctataaaaaa tttgtgaaaa aataatgttt attctgaaga tgtaaatttt gcaggaagat 720 tttattagaa tatggatcaa tatgcagtat tatgacctta tgatgaccta ttctttgaaa 780 agttqqqatt tactqtttta tacttaaacc ttttaaatqq ttttaaattc agatatqtaa 840 acaataggaa aaattgaaat tetteeaaaa atagtttaga ttattttgge ttattteaaa 900 atgtatcagt tottggtttt gtgatgttta tatttattat cttgacttca gt tac agg 958 ` Tyr Arg att tac caa gaa ttt gaa gta act gta cct gat tct atc act tct tgg 1006 Ile Tyr Gln Glu Phe Glu Val Thr Val Pro Asp Ser Ile Thr Ser Trp gtg gct act ggt ttt gtg atc tct gag gac ctg ggt ctt gga cta aca 1054 Val Ala Thr Gly Phe Val Ile Ser Glu Asp Leu Gly Leu Gly Leu Thr 20 30 act act cca gtg gag gtattgtatt aaagagctgc ttatcagtat tacggtgaca 1109

Thr Thr Pro Val Glu 35

	ttaagctaat	acagcgtcag	ctcctcaatt	tttttttaa	atgactgctt	ataatgttta	1169
	tcacagttta	gagattcctt	ggctttgtct	ttaggttttt	atctgtttta	tatttaagaa	1229
	tgtgagctat	atatagctat	ataaactgct	aaatgtgcaa	agtccgtatt	aagatttggg	1289
	tägaaaagtt	tattattgac	ctgaactaac	catctccaaa	ggccagaaga	gagagaaaga	1349
,	aaaagagaga	gagagaaaga	ggagaggaga	gagagagtga	gtctttctat	ttgtcctctt	1409
	caagaatgaa	cagaacttct	caagatgttc	cctagccaat	attccatcat	gtcttttggt	1469
,	caaattgcat	catatattgt	ttcctaagcc	agtcactggc	aggaggaata	taatgaccat	1529
•	gagtggcctg	aattttctca	tttgaaattg	aaatgtaatt	ttgatttaca	aaataatcgt	1589
	attcatgaaa	aatacagtgt	agattgaaaa	atgctttggg	tttatataga	aattggaatt	1649
i	agattgtaag	ctcaggccac	tataaacaga	caattcagca	acatgaatgt	ctgaagggac	1709
	attcaagaat	cattaggaac	atggggcaat	ttttcattgt	ctggggctgt	cctgagtatt	1769
	gcagactgtc	acccactaac	tacctatage	accttcgagt	catggtgaca	atctaagaca	1829
1	ccttcacaaa	tgtgcagata	aactctagag	ggagttactg	ctgccagcaa	aaccactggc	1889
,	ctaaactaac	ccaggtttag	ctttagatgc	aggtgtgggg	cttggccttt	tctgtaggac	1949
	ttggccaaca	atatcagaat	tgggtcactg	aggaggaagc	acatgtattc	agatgtccca	2009
•	cacattttct	catctgtatg	taaaaataaa	tcatatatat	gttttagaaa	taatttccaa	2069
	tttcctcttt	aaatttagtc	aggaagcaca	tgtattcaga	tgtcccacac	tagaacaggg	2129
•	gctgttggat	ttggcagggc	ttttaaagca	gattggtgga	gtcaatacag	catgaaagaa	2189
•	gagcaaattg	cttcgggatt	agacaggctg	ggttctagtt	ctggctcctc	tacttgccag	2249
•	caatatgaat	ttgtactagt	tacaaaaatc	tcaaaaattt	aattttcttt	tctataaact	2309
i	aggagactaa	cagtaacctt	atggggttgt	aataaccaaa	caaaataatt	tatgtgaagt	2369
•	gtttggttgc	tataaggcac	ttaataaagt	atagcaatta	ttatgttaag	taacataaat	2429
,	caagtcaatt	tgccgtcatt	catttgtgat	aagttgctgt	ttgctttctg	ttgatagcaa	2489
•	gttgacattt	ctagctgaag	ttaaaagctt	cacaggtttt	ataaagattg	catttaattg	2549
	cataaaatgt	gaagaatttt	gacctgaata	aaaatatgta	ctcgttgtgt	tctttccag	2608

<210> 6

<211> 23

<212> DNA

<213> HUMAN

wo	02/070738	PCT/CA02/00291
<400> ttcaaal	6 ttct tggtaaatcc tgt	23
<210>	7	
<211>	23	
<212>	DNA	
<213>	HUMAN .	•
	7 ttct tggtaaatcc tgg	23
<210>	8	
<211>	22	
	DNA	
<213>	HUMAN	
<400> atgacci	8 ttat gatgacctat tc	22
<210>	9	
<211>	18	
<212>	DNA .	
<213>	HUMAN	
<400> tttagai	9 ttat tttggctt	18
<210>	10	
<211>	18	
<212>	DNA	
<213>	HUMAN	
<400> atggtt	10 agtt caggtcaa	18
<210>	11	

<211> 27

<212>	DNA	
<213>	HUMAN	
<400> tgtatc	11 agtt cttggttttg tgatgtt	27
<210>	12	
<211>	30	
<212>	DNA	
<213>	HUMAN	
<400> ccaaga	12 agtg atagaatcag gtacagttac	. 30
<210>	13	
<211>	40	
<212>	DNA	
<213>	HUMAN .	
<400> tattate	13 cttg acttcagtta caggatttac caagaatttg	40
<210>	14	
<211>	37 [°]	
<212>	DNA .	
<213>	имам	
<400> tattato	14 cttg acttcagttc caggatttac caagaat	37
<210>	15	
<211>	18	
<212>	DNA	
<213>	synthetic construct	•

WO 02/070738

<220>

PCT/CA02/00291

PCT/CA02/00291

18

WO 02/070738 <221> primer <222> (1)..(18) <400> 15 gtagcccagg cagacgcc <210> 16 <211> 20 <212> DNA <213> synthetic construct

<220>

<221> primer

<222> (1)..(20)

<400> 16 20 cgcattgtta cactcttctc

<210> 17

<211> 19

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(19)

<400> 17 19 gattcttcaa atggacttt

<210> 18

<211> 21

<212> DNA

<213> synthetic construct

	·	
<220>		
<221>	primer	
<222>	(1)(21)	•
<400>	18	21
tgaatt	ccca atcctggagg a	21
<210>	19	
<211>	20	
<212>	DNA	
<213>	synthetic construct	
<220>		
<221>	primer	
<222>	(1)(20)	
<400>		
ttcaqq	aato togactetoo	20
ttcagg	aatg tggactctgg	20
<210>		20
	20	20
<210>	20	20
<210> <211> <212>	20	20
<210> <211> <212>	20 20 DNA	20
<210> <211> <212>	20 20 DNA	20
<210> <211> <212> <213> <220>	20 20 DNA	20
<210> <211> <212> <213> <223> <221> <222>	20 20 DNA synthetic construct	20
<210> <211> <212> <213> <220> <221>	20 20 DNA synthetic construct primer	20
<210> <211> <212> <213> <2213> <222> <400>	20 DNA synthetic construct primer (1)(20)	20
<210> <211> <212> <213> <2213> <222> <400>	20 DNA synthetic construct primer (1)(20)	
<210> <211> <212> <213> <223> <221> <222> <400> ctggga	20 DNA synthetic construct primer (1)(20)	

<212> DNA

<213> synthetic construct

<220>	•	
<221>	primer	
<222>	(1)(20)	
<400> acaatt	21 tcag acttctgagg	20
<210>	22	
<211>	21	
<212>	DNA	
<213>	synthetic construct	
<220>		
<221>	primer	
<222>	(1)(21)	
-		
<400>	22 gatc tatccaaaat c	21
JJ	· · · · · · · · · · · · · · · · · · ·	
<210>	23	
<211>	-	
<212>	DNA	
<213>	synthetic construct	
<220>		
	primer	
<222>	(1)(22)	
<400> gtgaca	23 acca ctgttggatc aa	22
.03.5		
<210>	24	
<211>	20	
<212>	DNA	

<213>	synthetic construct	٠	•
	·		

<220>

<221> primer

<222> (1)..(20)

<400> 24 tacatttctt gaaatacctg

20

<210> 25

<211> 18

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(18)

<400> 25 ggctgtgtca cagagatc

18

<210> 26

<211> 20

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(20)

<400> 26 gccacccaag aagtgataga

20

<210> 27

<211> 19

<212>	DNA	
<213>	synthetic construct	
<220>		
<221>	primer	

<400> 27

<222> (1)..(19)

cggcttcaag gaaacatct

<210> 28

<211> 22

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(22)

<400> 28 cagcaacatc taaatcaaag gc

<210> 29

<211> 18

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(18)

<400> 29 cacagccaaa gttccata

<210> 30

18

19 .

22

<211> 20

<212> DNA

<213> synthetic construct

<220>

<221> primer

<222> (1)..(20)

<400> 30 gctaggacct gttgtacacc

20

We claim:

1. An oligonucleotide comprising a sequence which binds specifically to (i) a region of CD109 nucleic acid that includes a single nucleotide polymorphism that is distinctive of a Gov^a allele and/or (ii) a region of CD109 nucleic acid that includes a single nucleotide polymorphism that is distinctive of a Gov^b allele.

- 2. The oligonucleotide of claim 1, comprising 8 to 50 nucleotides.
- 3. The oligonucleotide of claim 1, wherein the nucleic acid specifically binds to one of (i) or (ii) under high stringency hybridization conditions.
- 4. The oligonucleotide of claim 2, wherein the stringent hybridization conditions comprise 0.1XSSC, 0.1% SDS at 65°C.
- 5. The oligonucleotide of claim 1, wherein the CD109 nucleic acid comprises genomic DNA, cDNA, or RNA corresponding to the Gov^a allele of the CD109 gene or locus, or comprises genomic DNA, cDNA, or RNA corresponding to the Gov^b allele of the CD109 gene or locus.
- 6. The oligonucleotide of claim 5, wherein the Gov^a allele comprises an A at a position corresponding to position 2108 of SEQ ID NO:1 and corresponding to position 954 of SEQ ID NO:5.
- 7. The oligonucleotide of claim 5, wherein the Gov^b allele comprises a C at a position corresponding to position 2108 of SEQ ID NO:3 and corresponding to position 954 of SEQ ID NO:5.
- 8. The oligonucleotide of claim 1, comprising a sequence complementary to the Gov^a allele or to the Gov^b allele.
- 9. The oligonucleotide of claim 6, comprising a sequence selected from the group consisting of:
- (a) 8-50 nucleotides of SEQ ID NO:1;
- (b) a sequence that is complementary to a sequence specified in (a); and
- (c) a sequence having at least 70% sequence identity to a sequence in (a) or (b), wherein the sequence having identity is capable of hybridization to CD109 under high stringency hybridization conditions.

10. The oligonucleotide of claim 7, comprising a sequence selected from the group consisting of:

- (a) 8-50 nucleotides of SEQ ID NO:3;
- (b) a sequence that is complementary to a sequence specified in (a); and
- (c) a sequence having at least 70% sequence identity to a sequence in (a) or (b), wherein the sequence having identity is capable of hybridization to CD109 under high stringency hybridization conditions.
- 11. A oligonucleotide comprising all or part of any one of SEQ ID NO:6-SEQ ID NO:14 or a complement thereof.
- 12. The oligonucleotide of claim 11, comprising 8 to 50 nucleic acids.
- 13. The oligonucleotide of claim 1, wherein the nucleic acid is capable of use as a probe in a hybridization assay.
- 14. The oligonucleotide of claim 1, wherein the nucleic acid sequence is detectably labelled.
- 15. The oligonucleotide of claim 14, wherein the detectable label comprises:
- (a) a fluorogenic dye; and/or
- (b) a biotinylation modification; and/or
- (c) a radiolabel.
- 16. The oligonucleotide of claim 1, wherein the sequence comprises DNA, a DNA analog, RNA or an RNA analog.
- 17. The oligonucleotide of claim 1, wherein the oligonucleotide is attached to a substrate.
- 18. The oligonucleotide of claim 1, wherein the oligonucleotide is capable of use as a primer that will specifically bind proximate to, and/or cause elongation through, a CD109 sequence, including the single nucleotide polymorphism distinctive of the Gov^a or Gov^b alleles.
- 19. A Gov genotyping kit comprising a detection agent for detecting the presence of a Gov allelespecific target sequence in a CD109 nucleic acid derived from a subject.

20. The kit of claim 19, wherein the detection agent comprises a nucleic acid and/or a restriction enzyme.

- 21. The kit of claim 19, further comprising a container.
- 22. The kit of claim 21, wherein the container comprises a biological sample container for housing the detection agent.
- 23. The kit of claim 19, further comprising a plate having a plurality of wells and having bound thereto probes having a nucleic acid sequence which specifically binds to a CD109 sequence including a Gov^a or a Gov^b allele target sequence.
- 24. The kit of claim 20, wherein the restriction enzyme is selected from the group consisting of Bst2UI, BstNI, BstOI, EcoRII, MaeIII, MspR91, MvaI, ScrFI or an isoschizomer thereof.
- 25. The kit of claim 19, further comprising an amplification agent for amplifying the nucleic acid.
- 26. The kit of claim 25, wherein the amplification agent amplifies a region of CD109 platelet, T cell, or endothelial cell mRNA including the single nucleotide polymorphism distinctive of a Gov^a or Gov^b allele.
- 27. The kit of claim 25, wherein the amplification agent comprises a primer set including first and second primers, wherein the first primer is a nucleic acid that will specifically bind proximate to, and/or cause elongation through, CD109 sequence that includes the single nucleotide polymorphism distinctive of a Gov^a allele and the second primer is a nucleic acid that will specifically bind proximate to, and/or cause elongation through, CD109 sequence that includes the single nucleotide polymorphism distinctive of a Gov^b allele.
- 28. The kit of claim 19, wherein the nucleic acid is obtained by amplification with all or part of the nucleic acid of any one of SEQ ID NO:6 SEQ ID NO:14 or the complement thereof.
- 29. The kit of claim 19, further comprising all or part of a CD109 gene, a CD109-encoding mRNA, or a CD109 cDNA made from a CD109-encoding mRNA.
- 30. The kit of claim 19, comprising the oligonucleotide of any of claims 1 to 18.

31. The kit of claim 19, for detecting that the subject has or is at risk of a disease, disorder or abnormal physical state.

- 32. The kit of claim 31, wherein the disease, disorder or abnormal physical state comprises a blood disease, disorder or abnormal physical state.
- 33. The kit of claim 32, wherein the blood disease, disorder or abnormal physical state comprises bleeding of the subject, or increased risk of bleeding, due to destruction of blood platelets.
- 34. The kit of claim 33, wherein the blood disease, disorder or abnormal physical state comprises post-transfusion purpura ("PTP"), post-transfusion platelet refractoriness ("PR") or neonatal alloimmune thrombocytopenia ("NAIT").
- 35. The kit of claim 33, wherein the nucleic acid is obtained from mRNA from human platelets, T cells, endothelial cells, or human genomic DNA.
- 36. A method of Gov alloantigen genotyping a subject comprising:
- (a) providing a CD109 nucleic acid sample derived from the subject; and
- (b) detecting a region of CD109 nucleic acid that includes a single nucleotide polymorphism distinctive of a Gov^a or a Gov^b allele.
- 37. The method of claim 36 comprising determining whether the subject is homozygous or heterozygous for the Gov alleles.
- 38. The method of claim 37, wherein the subject is a human and the Gov genotype is used to determine that the subject has, or is at risk of a disease, disorder or abnormal physical state.
- 39. The method of claim 38, wherein the disease, disorder or abnormal physical state comprises a blood disease, disorder or abnormal physical state.
- 40. The method of claim 39, wherein the blood disease, disorder or abnormal physical state comprises bleeding of the subject, or increased risk of bleeding, due to destruction of blood platelets.

41. The method of claim 40, wherein the blood disease, disorder or abnormal physical state comprises post-transfusion purpura ("PTP"), post-transfusion platelet refractoriness ("PR") or neonatal alloimmune thrombocytopenia ("NAIT").

- 42. The method of claim 41, wherein the nucleic acid is obtained by amplifying the nucleic acid from the subject.
- 43. The method of claim 42, wherein the nucleic acid is obtained by amplification with all or part of the oligonucleotide of any of claims 1 to 18.
- 44. The method of claim 41, wherein the nucleic acid is obtained from mRNA from human platelets, T cells, endothelial cells, or human genomic DNA.
- 45. The method of claim 36, wherein the detection step comprises determining the nucleotide sequence of the CD109 nucleic acid.
- 46. The method of clam 36, wherein the detection step comprises contacting the nucleic acid with the oligonucleotide of any of claims 1 to 18 under high stringency conditions.
- 47. The method of claim 46, wherein the oligonucleotide will selectively hybridize to (i) a region of CD109 nucleic acid that includes a single polymorphism distinctive of a Gov^a allele or (ii) a region of CD109 nucleic acid that includes a single polymorphism distinctive of a Gov^b allele.
- 48. The method of claim 36, wherein the detecting step comprises:
- (a) performing a restriction endonuclease digestion of the nucleic acid, thereby providing a nucleic acid digest; and
- (b) contacting the digest with the oligonucleotide.
- 49. The method of claim 47, wherein the hybridization occurs either during or subsequent to PCR amplification and the analysis is by "Real-Time" PCR analysis, or fluorimetric analysis.
- 50. The method of claim 36, wherein the detection step comprises:
- (a) incubation of the amplified nucleic acid with a restriction endonuclease under conditions whereby the DNA will be cleaved if the nucleic acid comprises a recognition site for the enzyme; and

(b) determining whether the nucleic acid contains a recognition site for the restriction enzyme characteristic of cDNA made from mRNA encoding a Gov^a or Gov^b allele of CD109.

- 51. The method of claim 50, wherein the restriction enzyme is selected from the group consisting of Bst2UI, BstNI, BstOI, EcoRII, MaeIII, MspR91, MvaI, ScrFI or an isoschizomer thereof.
- 52. The method of claim 50, wherein the determination step includes size analysis of the nucleic acid.
- 53. The method of claim 50, wherein the amplified nucleic acid is analyzed by electrophoretic mobility and the mobility of the amplified nucleic acid is compared to the characteristic mobility of amplified nucleic acid fragments corresponding to the Gov^a or Gov^b alleles of CD109.
- 54. A method of amplifying CD109 mRNA comprising amplifying the mRNA by PCR using an oligonucleotide of any of claims 1 to 18.
- 55. A Gov^a specific antibody.
- 56. The antibody of claim 55, that recognizes specifically a Gov^a allele-specific CD109 epitope corresponding to the polypeptide encoded by a CD109 nucleic acid containing an A at the position corresponding to position 2108 of SEQ ID NO:1 and position 954 of SEQ ID NO:5., and containing the amino acid Tyrosine at the position corresponding to position 703 of the CD109 protein encoded by SEQ ID NO:1.
- 57. A Gov^b specific antibody.
- 58. The antibody of claim 57, that recognizes specifically a Gov^b allele-specific CD109 epitope corresponding to the polypeptide encoded by a CD109 nucleic acid containing a C at the position corresponding to position 2108 of SEQ ID NO:3 and position 954 of SEQ ID NO:5., and containing the amino acid Serine at the position corresponding to position 703 of the CD109 protein encoded by SEQ ID NO:3.
- 59. The antibody of any of claims 55 to 58, further comprising a monoclonal antibody or a polyclonal antibody.
- 60. The antibody of any of claims 55 to 59, further comprising a detectable label.

61. An immunogenic composition comprising a Gov specific antibody of any of claims 55 to 60.

- 62. A method of Gov alloantigen phenotyping a subject, comprising:
- (a) providing a CD109 polypeptide sample derived from the subject; and
- (b) detecting the presence of a Gov^a or a Gov^b antigen in the CD109 polypeptide.
- 63. The method of claim 62, wherein the CD109 is membrane bound CD109 or isolated CD109.
- 64. The method of claim 62, wherein the detection step comprises contacting the polypeptide sample with the antibody of any of claims 55 to 60.
- 65. A diagnostic kit for Gov alloantigen phenotyping a subject, comprising a Gov^a antibody and/or a Gov^b antibody of any of claims 55 to 60.
- 66. The kit of claim 65, further comprising a container.
- 67. An isolated polypeptide containing Gov^a allele-specific amino acid sequence and which is specifically reactive with a Gov^a antibody.
- 68. An isolated polypeptide containing Gov^b allele-specific amino acid sequence and which is specifically reactive with a Gov^b antibody.
- 69. The isolated polypeptide of claim 67 or 68, comprising between 4 and 100 amino acids.
- 70. The isolated polypeptide of claim 67, 68, or 69, comprising a full-length CD109 polypeptide, or a fragment of a CD109 polypeptide.
- 71. An isolated CD109 polypeptide fragment, comprising a Gov^a or a Gov^b antigen.
- 72. The polypeptide fragment of claim 71, comprising all of, or a fragment of, the protein encoded by SEQ ID NO:1., and in which the amino acid corresponding to position 703 of the protein encoded by SEQ ID NO:1 is a Tyrosine

73. The polypeptide fragment of claim 72, comprising all of, or a fragment of, the protein encoded by SEQ ID NO:3., and in which the amino acid corresponding to position 703 of the protein encoded by SEQ ID NO:3 is a Serine.

- 74. The polypeptide fragment of claim 72 or 73, comprising between 4 and 100 amino acids.
- 75. The polypeptide fragment of claim 74, comprising between 7 and 50 amino acids.
- 76. The polypeptide fragment of claims 67 to 75, in which the polypeptide is purified from native CD109, is synthetic, or is prepared by recombinant means.
- 77. The polypeptide fragment of claims 67 to 76, in which the polypeptide is bound to a substrate.
- 78. A fusion compound comprising the polypeptide of any of claims 67 to 77 connected to an immunogenic carrier.
- 79. The fusion compound of claim 78, wherein the immunogenic carrier comprises a proteinaceous carrier.
- 80. The fusion compound of claim 79, wherein the immunogenic carrier comprises a detectable label.
- 81. A Gov^a or Gov^b specific antibody recognizing the fusion compound of claim 80.
- 82. An immunogenic composition comprising the polypeptide, polypeptide fragment or fusion compound of any of claims 67 to 80.
- 83. A method of producing a Gov^a or Gov^b specific antibody, comprising contacting an animal with the immunogenic composition of claim 82, so that the animal produces antibodies against the immunogenic composition.
- 84. The method of claim 83, wherein the animal is a bird or a mammal.
- 85. A method of screening an antibody producing culture to determine whether the culture produces Gov^a or Gov^b specific antibody, comprising:
- (a) contacting a polypeptide of the invention with the culture; and

- (b) detecting Gov^a or Gov^b specific antibody.
- 86. The method of claim 85, wherein the polypeptide comprises a detectable label.
- 87. The method of claim 86, wherein the polypeptide is attached to a substrate.
- 88. A method of purifying a Gov allele-specific antibody from a sample, comprising:
- (a) contacting a Gov allele-specific antibody with a polypeptide of the invention comprising a Gov^a or Gov^b antigen, so that an antibody:polypeptide complex is formed;
- (b) separating the complex from the sample; and
- (c) next separating the antibody from the polypeptide.
- 89. The method of claim 88, wherein the polypeptide is bound to a substrate.
- 90. The method of claim 89, wherein the polypeptide comprises a detectable label.
- 91. A method of purifying a Gov polypeptide from a sample, comprising:
- (a) contacting a Gov allele-specific antibody with a polypeptide of the invention containing a Gov^a or Gov^b-specific epitope, so that an antibody:polypeptide complex is formed;
- (b) separating the complex from the sample; and
- (c) next separating the antibody from the polypeptide.
- 92. The method of claim 91, wherein the antibody is bound to a substrate.
- 93. The method of claim 92, wherein the antibody comprises a detectable label.
- 94. A method of screening a subject sample to determine whether the sample contains Gov^a or Gov^b -specific antibodies, comprising:
- (a) contacting a polypeptide of the invention with the sample; and
- (b) detecting the presence or absence of Gov^a or Gov^b specific antibody.
- 95. The method of claim 94, wherein the polypeptide comprises a detectable label.
- 96. The method of claim 95, wherein the polypeptide is attached to a substrate

97. The method of claim 96, wherein the subject comprises a mother of a fetus or a newborn infant, or the fetus or newborn infant itself, and the presence of Gov^b -specific antibody indicates that the fetus or infant has, or is at risk of NAIT.

- 98. The method of claim 97, wherein the presence of Gov^a or Gov^b specific antibody indicates that the subject has, or is at risk of a blood disease, disorder or abnormal physical state.
- 99. The method of claim 98, wherein the blood disease, disorder or abnormal physical state comprises bleeding of the subject, or increased risk of bleeding, due to destruction of blood platelets.
- 100. The method of claim 99, wherein the blood disease, disorder or abnormal physical state comprises post-transfusion purpura ("PTP"), post-transfusion platelet refractoriness ("PR") or neonatal alloimmune thrombocytopenia (NAIT).
- 101. The method of claim 100, wherein the sample comprises human serum or plasma.
- 102. A diagnostic kit for detection of Gov^a or Gov^b specific antibody, comprising a polypeptide of any of claims 67 to 80.
- 103. The kit of claim 102, further comprising a container.
- 104. A method of determining Gov antibody specificity, comprising:
- (a) contacting an antibody with a first polypeptide comprising a Gov^a antigen and a second polypeptide comprising a Gov^b antigen; and
- (b) determining whether the antibody binds to either or both of the first and second polypeptide.
- 105. A method of blocking Gov^a antibody binding to an antigen, comprising: contacting the antibody with a polypeptide of the invention comprising a Gov^a antigen so that an antibody:polypeptide complex is formed.
- 106. The method of claim 105, wherein the polypeptide comprises a detectable label.
- 107. The method of claim 106, wherein the polypeptide is bound to a substrate.

108. A method of blocking Gov^b antibody binding to an antigen, comprising: contacting the antibody with a polypeptide of the invention comprising a Gov^b antigen so that an antibody:polypeptide complex is formed.

- 109. The method of claim 108, wherein the polypeptide comprises a detectable label.
- 110. The method of claim 109, wherein the polypeptide is bound to a substrate.
- 111. A pharmaceutical composition comprising the polypeptide of any of claims 67 to 80.
- 112. A method of immunizing a subject so that the subject will produce anti-idiotypic antibodies, comprising administering to the subject the immunogenic composition of any of claims 82 to 84.
- 113. A method of blocking Gov^a or Gov^b specific antibodies from binding to CD109 in a subject, comprising: administering to the subject polypeptides of the invention capable of binding to Gov^a and/or Gov^b specific antibodies.
- 114. The method of claim 113, wherein the polypeptide comprises a detectable label.
- 115. The method of claim 114, wherein the binding of the polypeptide to the Gov^a or Gov^b specific antibody prevents alloimmune cell destruction by the antibody.
- 116. The method of claim 115, wherein the binding of the polypeptide to the Gov^a or Gov^b specific antibody depletes the antibody.
- 117. The method of claim 116, wherein the subject has or is at risk of a blood disease, disorder or abnormal physical state.
- 118. The method of claim 117, wherein the blood disease, disorder or abnormal physical state comprises bleeding of the subject, or increased risk of bleeding, due to alloimmune destruction of blood platelets.
- 119. The method of claim 118, wherein the blood disease, disorder or abnormal physical state comprises post-transfusion purpura ("PTP"), post-transfusion platelet refractoriness ("PR") or neonatal alloimmune thrombocytopenia ("NAIT").